

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 7/48, 7/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/52530</b> <b>(43) International Publication Date:</b> 26 November 1998 (26.11.98)
<b>(21) International Application Number:</b> PCT/IB98/00783 <b>(22) International Filing Date:</b> 20 May 1998 (20.05.98)  <b>(30) Priority Data:</b> 08/862,774      23 May 1997 (23.05.97)      US  <b>(71) Applicant:</b> THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).  <b>(72) Inventors:</b> ZUKOWSKI, Joseph, Michael; 3811 Drakewood Drive, Cincinnati, OH 45209 (US). DECKNER, George, Endel; 10572 Tanager Hills Drive, Cincinnati, OH 45249 (US). SaNOGUEIRA, James, Pedrosa, Jr.; 4 Kimmissy Court, Suffern, NY 10901 (US).  <b>(74) Agents:</b> REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		<b>(81) Designated States:</b> AU, CA, CN, CZ, JP, KR, MX, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> SKIN CARE COMPOSITIONS COMPRISING VITAMIN B3 AND A PRESERVATIVE  <b>(57) Abstract</b>  Disclosed are skin care compositions containing a vitamin B <sub>3</sub> compound which generally improve the quality of the skin, particularly human facial skin. More particularly, the present invention relates to niacinamide containing skin care compositions with improved skin compatibility.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

SKIN CARE COMPOSITIONS COMPRISING VITAMIN B<sub>3</sub> AND A PRESERVATIVETECHNICAL FIELD

5 Invention relates to skin care compositions containing a vitamin B<sub>3</sub> compound which generally improve the quality of the skin, particularly human facial skin. More particularly, the present invention relates to niacinamide containing skin care compositions with improved skin compatibility.

BACKGROUND OF THE INVENTION

10 Many personal care products currently available to consumers are directed primarily to improving the health and/or physical appearance of the skin. Among these skin care products, many are directed to delaying, minimizing or even eliminating skin wrinkling and other histological changes typically associated with the aging of skin or environmental damage to human skin.

15 Skin is subject to insults by many extrinsic and intrinsic factors. Extrinsic factors include ultraviolet radiation (e.g., from sun exposure), environmental pollution, wind, heat, low humidity, harsh surfactants, abrasives, and the like. Intrinsic factors include chronological aging and other biochemical changes from within the skin. Whether extrinsic or intrinsic, these factors result in visible signs of  
20 skin aging and environmental damage, such as wrinkling and other forms of roughness (including increased pore size, flaking and skin lines), and other histological changes associated with skin aging or damage. To many people, skin wrinkles are a reminder of the disappearance of youth. As a result, the elimination of wrinkles has become a booming business in youth-conscious societies.  
25 Treatments range from cosmetic creams and moisturizers to various forms of cosmetic surgery.

Extrinsic or intrinsic factors may result in the thinning and general degradation of the skin. For example, as the skin naturally ages, there is a reduction in the cells and blood vessels that supply the skin. There is also a flattening of the  
30 dermal-epidermal junction which results in weaker mechanical resistance of this junction. See, for example, Oikarinen, "The Aging of Skin: Chronoaging Versus Photoaging," *Photodermatol. Photoimmunol. Photomed.*, vol. 7, pp. 3-4, 1990, which is incorporated by reference herein in its entirety.

Vitamin B<sub>3</sub> compounds, particularly niacinamide, have recently been found  
35 to provide measurable skin regulating benefits. For example, topical niacinamide helps to regulate the signs of skin aging, i.e., reduce or efface the visibility of the fine lines, wrinkles, and other forms of uneven or rough surface texture associated

with aged or photodamaged skin. These compounds have also been found useful in reducing the overall oiliness of skin.

In formulating products containing vitamin B<sub>3</sub> compounds, much attention is directed toward providing compositions which deliver and retain optimal concentrations of the vitamin B<sub>3</sub> compounds in the stratum corneum with minimum absorption into the systemic circulation. Furthermore, promoting user compliance with respect to chronic treatment regimens is also important. Current vitamin B<sub>3</sub> formulations, however, can be drying and irritating. Such formulations may cause individuals to refrain from using vitamin B<sub>3</sub> products as frequently and copiously as is necessary for optimum benefit.

The present inventors have found that compositions containing natural or synthetic vitamin B<sub>3</sub> compounds along with a preservative component comprising a formaldehyde donating preservative and a halopropynyl compound, deliver the skin regulating benefits of a vitamin B<sub>3</sub> compound with reduced dryness and/or irritation. These compositions have improved user acceptance and, thus, promote better user compliance with a concomitant overall improvement in skin regulating benefit.

It is, therefore, an object of the present invention is to provide natural or synthetic vitamin B<sub>3</sub> containing skin care compositions having improved skin compatibility.

Another object of the present invention is to provide natural or synthetic vitamin B<sub>3</sub> compositions containing preservative systems which provide preservation activity at concentrations of no more than 0.2%.

Still another object of the present invention is to provide natural or synthetic vitamin B<sub>3</sub> compositions containing preservative systems which do not substantially impact niacinamide stability or bioavailability.

These and other objects of this invention will become apparent in light of the following disclosure.

### SUMMARY OF THE INVENTION

The present invention relates to skin care compositions, comprising:

- a.) from about 0.01% to about 50% of a vitamin B<sub>3</sub> compound, and
- b.) from about 0.001% to about 5% of a preservative component, comprising:
  - i.) a formaldehyde donor; and
  - ii.) a halopropynyl compound selected from the group consisting of iodopropargyl esters, ethers, acetals, carbamates and carbonates.

The present invention further relates to methods of regulating skin conditioning.

### DETAILED DESCRIPTION OF THE INVENTION

All percentages and ratios used herein are by weight of the total composition and all measurements made are at 25°C. unless otherwise designated.

5       The compositions of the present invention can comprise, consist essentially of, or consist of, the essential as well as optional ingredients and components described herein. As used herein, "consisting essentially of" means that the composition or component may include additional ingredients, but only if the additional ingredients do not materially alter the basic and novel characteristics of  
10       the claimed compositions or methods.

All publications cited herein are hereby incorporated by reference in their entirety.

The term "topical application", as used herein, means to apply or spread the compositions of the present invention onto the surface of the skin.

15       The term "dermatologically-acceptable," as used herein, means that the compositions or components thereof so described are suitable for use in contact with human skin without undue toxicity, incompatibility, instability, allergic response, and the like.

20       The term "safe and effective amount" as used herein means an amount of a compound or composition sufficient to significantly induce a positive benefit, preferably a positive skin appearance or feel benefit, including independently the benefits disclosed herein, but low enough to avoid serious side effects, i.e., to provide a reasonable benefit to risk ratio, within the scope of sound judgment of the skilled artisan.

25       The term "skin compatibility," as used herein means the ability of skin to tolerate long term application of topical compositions with minimal adverse skin reactions such as stinging, burning, redness, itching and folliculitis.

30       The compositions of the present invention are useful for topical application and for regulating skin condition, including visible and/or tactile discontinuities in skin (especially the skin surface; such discontinuities are generally undesirable). Such discontinuities may be induced or caused by internal and/or external factors, and include the signs of skin aging described herein. The term "regulating skin condition" includes prophylactically regulating and/or therapeutically regulating skin condition, including visible and/or tactile discontinuities in skin. As used herein,  
35       prophylactically regulating skin condition includes delaying, minimizing and/or preventing visible and/or tactile discontinuities in skin. As used herein, therapeutically regulating skin condition includes ameliorating, e.g., diminishing,

minimizing and/or effacing, discontinuities in skin. Regulating skin condition involves improving skin appearance and/or feel.

The compositions of the present invention are useful for regulating signs of skin aging, more especially visible and/or tactile discontinuities in skin texture associated with aging. "Regulating the signs of skin aging" includes prophylactically regulating and/or therapeutically regulating one or more of such signs (similarly, regulating a given sign of skin aging, e.g., lines, wrinkles or pores, includes prophylactically regulating and/or therapeutically regulating that sign). As used herein, prophylactically regulating such signs includes delaying, minimizing and/or preventing signs of skin aging. As used herein, therapeutically regulating such signs includes ameliorating, e.g., diminishing, minimizing and/or effacing signs of skin aging.

"Signs of skin aging" include, but are not limited to, all outward visibly and tactilely perceptible manifestations as well as any other macro or micro effects due to skin aging. Such signs may be induced or caused by intrinsic factors or extrinsic factors, e.g., chronological aging and/or environmental damage. These signs may result from processes which include, but are not limited to, the development of textural discontinuities such as wrinkles, including both fine superficial wrinkles and coarse deep wrinkles, skin lines, crevices, bumps, large pores (e.g., associated with adnexal structures such as sweat gland ducts, sebaceous glands, or hair follicles), scaliness, flakiness and/or other forms of skin unevenness or roughness, loss of skin elasticity (loss and/or inactivation of functional skin elastin), sagging (including puffiness in the eye area and jowls), loss of skin firmness, loss of skin tightness, loss of skin recoil from deformation, discoloration (including under-eye circles), blotching, sallowness, hyperpigmented skin regions such as age spots and freckles, keratoses, abnormal differentiation, hyperkeratinization, elastosis, collagen breakdown, and other histological changes in the stratum corneum, dermis, epidermis, the skin vascular system (e.g., telangiectasia or spider vessels), and underlying tissues, especially those proximate to the skin.

It is to be understood that the present invention is not to be limited to regulation of the above mentioned "signs of skin aging" which arise due to mechanisms associated with skin aging, but is intended to include regulation of said signs irrespective of the mechanism of origin. As used herein, "regulating skin condition" is intended to include regulation of such signs irrespective of the mechanism of origin.

The present invention is especially useful for therapeutically regulating visible and/or tactile discontinuities in mammalian skin texture, including texture

discontinuities associated with skin aging. As used herein, therapeutically regulating such discontinuities includes ameliorating, e.g., diminishing, minimizing and/or effacing visible and/or tactile discontinuities in the texture of mammalian skin, to thereby provide improved skin appearance and/or feel, e.g., a smoother, more even appearance and/or feel. Such visible and/or tactile discontinuities in skin texture include crevices, bumps, pores, fine lines, wrinkles, scales, flakes and/or other forms of textural unevenness or roughness associated with skin aging. For example, the length, depth, and/or other dimension of lines and/or wrinkles are decreased, the apparent diameter of pores decreases, or the apparent height of tissue immediately proximate to pore openings approaches that of the interadnexal skin.


The present invention is also especially useful for prophylactically regulating visible and/or tactile discontinuities in mammalian skin texture, including texture discontinuities associated with skin aging. As used herein, prophylactically regulating such discontinuities includes delaying, minimizing and/or preventing visible and/or tactile discontinuities in the texture of mammalian skin, to thereby provide improved skin appearance and/or feel, e.g., a smoother, more even appearance and/or feel.

### Essential Components

#### Vitamin B<sub>3</sub> component

The compositions of the present invention comprise as a safe and effective amount of a natural or synthetic vitamin B<sub>3</sub> compound. The compositions of the present invention preferably comprise from about 0.01% to about 50%, more preferably from about 0.1% to about 10%, even more preferably from about 0.5% to about 10%, and still more preferably from about 1% to about 5%, most preferably from about 2% to about 5%, of the vitamin B<sub>3</sub> compound.

As used herein, "vitamin B<sub>3</sub> compound" means a compound having the formula:




wherein R is - CONH<sub>2</sub> (i.e., niacinamide), - COOH (i.e., nicotinic acid) or - CH<sub>2</sub>OH (i.e., nicotinyl alcohol); derivatives thereof; and salts of any of the foregoing.

Exemplary derivatives of the foregoing vitamin B<sub>3</sub> compounds include nicotinic acid esters, including non-vasodilating esters of nicotinic acid, nicotinyl amino acids, nicotinyl alcohol esters of carboxylic acids, nicotinic acid N-oxide and niacinamide N-oxide.


Suitable esters of nicotinic acid include nicotinic acid esters of C<sub>1</sub>-C<sub>22</sub>, preferably C<sub>1</sub>-C<sub>16</sub>, more preferably C<sub>1</sub>-C<sub>6</sub> alcohols. The alcohols are suitably straight-chain or branched chain, cyclic or acyclic, saturated or unsaturated (including aromatic), and substituted or unsubstituted. The esters are preferably non-rubicient. As used herein, "non-rubicient" means that the ester does not commonly yield a visible flushing response after application to the skin in the subject compositions (the majority of the general population would not experience a visible flushing response, although such compounds may cause vasodilation not visible to the naked eye). Non-rubicient esters of nicotinic acid include tocopherol nicotinate and inositol hexanicotinate; tocopherol nicotinate is preferred.

Other derivatives of the vitamin B<sub>3</sub> compound are derivatives of niacinamide resulting from substitution of one or more of the amide group hydrogens. Nonlimiting examples of derivatives of niacinamide useful herein include nicotinyl amino acids, derived, for example, from the reaction of an activated nicotinic acid compound (e.g., nicotinic acid azide or nicotinyl chloride) with an amino acid, and nicotinyl alcohol esters of organic carboxylic acids (e.g., C<sub>1</sub> - C<sub>18</sub>). Specific examples of such derivatives include nicotinuric acid and nicotinyl hydroxamic acid, which have the following chemical structures:

nicotinuric acid:



nicotinyl hydroxamic acid:



Exemplary nicotinyl alcohol esters include nicotinyl alcohol esters of the carboxylic acids salicylic acid, acetic acid, glycolic acid, palmitic acid and the like. Other non-limiting examples of vitamin B<sub>3</sub> compounds useful herein are 2-chloronicotinamide, 6-aminonicotinamide, 6-methylnicotinamide, n-methylnicotinamide, n,n-diethylnicotinamide, n-(hydroxymethyl)-nicotinamide, quinolinic acid imide, nicotinanilide, n-benzylnicotinamide, n-ethylnicotinamide, nifenazone.



nicotinaldehyde, isonicotinic acid, methyl isonicotinic acid, thionicotinamide, nialamide, 1-(3-pyridylmethyl) urea, 2-mercaptonicotinic acid, nicomol, and niaprazine.

Examples of the above vitamin B<sub>3</sub> compounds are well known in the art and  
5 are commercially available from a number of sources, e.g., the Sigma Chemical Company (St. Louis, MO); ICN Biomedicals, Inc. (Irvin, CA) and Aldrich Chemical Company (Milwaukee, WI).

One or more vitamin B<sub>3</sub> compounds may be used herein. Preferred vitamin B<sub>3</sub> compounds are niacinamide and tocopherol nicotinate. Niacinamide is more  
10 preferred.

When used, salts, derivatives, and salt derivatives of niacinamide are preferably those having substantially the same efficacy as niacinamide in the methods of regulating skin condition described herein.

Salts of the vitamin B<sub>3</sub> compound are also useful herein. Nonlimiting  
15 examples of salts of the vitamin B<sub>3</sub> compound useful herein include organic or inorganic salts, such as inorganic salts with anionic inorganic species (e.g., chloride, bromide, iodide, carbonate, preferably chloride), and organic carboxylic acid salts (including mono-, di- and tri- C1 - C18 carboxylic acid salts, e.g., acetate, salicylate, glycolate, lactate, malate, citrate, preferably monocarboxylic acid salts such as  
20 acetate). These and other salts of the vitamin B<sub>3</sub> compound can be readily prepared by the skilled artisan, for example, as described by W. Wenner, "The Reaction of L-Ascorbic and D-Isoascorbic Acid with Nicotinic Acid and Its Amide", J. Organic Chemistry, VOL. 14, 22-26 (1949), which is incorporated herein by reference. Wenner describes the synthesis of the ascorbic acid salt of niacinamide.

25 In a preferred embodiment, the ring nitrogen of the vitamin B<sub>3</sub> compound is substantially chemically free (e.g., unbound and/or unhindered), or after delivery to the skin becomes substantially chemically free ("chemically free" is hereinafter alternatively referred to as "uncomplexed"). More preferably, the vitamin B<sub>3</sub> compound is essentially uncomplexed. Therefore, if the composition contains the  
30 vitamin B<sub>3</sub> compound in a salt or otherwise complexed form, such complex is preferably substantially reversible, more preferably essentially reversible, upon delivery of the composition to the skin. For example, such complex should be substantially reversible at a pH of from about 5.0 to about 6.0. Such reversibility can be readily determined by one having ordinary skill in the art.

35 More preferably the vitamin B<sub>3</sub> compound is substantially uncomplexed in the composition prior to delivery to the skin. Exemplary approaches to minimizing or preventing the formation of undesirable complexes include omission of materials

which form substantially irreversible or other complexes with the vitamin B<sub>3</sub> compound, pH adjustment, ionic strength adjustment, the use of surfactants, and formulating wherein the vitamin B<sub>3</sub> compound and materials which complex therewith are in different phases. Such approaches are well within the level of  
5 ordinary skill in the art.

Thus, in a preferred embodiment, the vitamin B<sub>3</sub> compound contains a limited amount of the salt form and is more preferably substantially free of salts of a vitamin B<sub>3</sub> compound. Preferably the vitamin B<sub>3</sub> compound contains less than about 50% of such salt, and is more preferably essentially free of the salt form. The  
10 vitamin B<sub>3</sub> compound in the compositions hereof having a pH of from about 4 to about 7 typically contain less than about 50% of the salt form.

The vitamin B<sub>3</sub> compound may be included as the substantially pure material, or as an extract obtained by suitable physical and/or chemical isolation from natural (e.g., plant) sources. The vitamin B<sub>3</sub> compound is preferably  
15 substantially pure, more preferably essentially pure.


#### Preservative Component

Also essential to the compositions of the present invention is a preservative component comprising a formaldehyde donor and a halopropynyl compound. The preservative component of the present invention provides substantially no negative  
20 impact to the active's stability. The phrase, "substantially no negative impact," as used herein, means that less than about 10% of the active is chemically degraded or otherwise altered resulting in an inactive degradation product when in the presence of the preservative component in the composition herein during a one month period at a temperature of 40°C.

25 Suitable formaldehyde donors have chemical and physical characteristics compatible with use in personal care products. Such products preferably are not odiferous or an irritant or toxic when applied to the skin. Examples of suitable formaldehyde donors include dimethyloldimethylhydantoin, N,N"-methylene bis [N'-(hydroxymethyl)-2,5-dioxo-4-imidazolidinyl]urea; N-(hydroxymethyl)-N-(1,3-  
30 dihydroxymethyl-2,5-dioxo-4-imidazolidinyl)-N'-(hydroxymethyl) urea; the cis isomer of 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride, sodium hydroxymethylglycinate, dimethyl oxazoline 7-ethylbicyclohexasolidine, 2-bromo-2-nitropropane-1,3-diol, 5-bromo-5-nitro-1,3-dioxane and mixtures thereof.

Examples of suitable halopropynyl compounds include compounds derived  
35 from propargyl or iodopropargyl alcohols such as the esters, ethers, acetals, carbamates and carbonates and the iodopropargyl derivatives of pyrimidines, triazolinones, tetrazoles, triazinones, sulfamides, benzothiazoles, ammonium salts.

carboxamides, hydroxamates, ureas and mixtures thereof. Preferred among these compounds is 3-iodo-2-propynylbutyl carbamate (IPBC). This compound is included within the broadly useful class of compounds having the generic formula:



wherein R is selected from the group consisting of substituted and unsubstituted alkyl, aryl and alkylaryl groups having from 1 to 20 carbon atoms, and m and n are independent integers from 1 to 3.

The weight ratio of the two components varies depending on the application and the particular component selected. Broadly speaking, the weight ratio varies from about 2000:1 to about 1:1 parts of the first to the second component may be used, preferably from about 500:1 to about 1:1, most preferably from about 200:1 to about 1:1, the parts being by weight. The preservative component of the present invention is present at a concentration of from about 0.001% to about 5%, preferably from about 0.01% to about 3%, more preferably from about 0.01% to about 2%, more preferably from about 0.05% to about 1% and most preferably from about 0.05% to about 0.2%.

#### Optional Components

##### Carrier

The compositions of the present invention comprise from about 1% to about 99.5% of a dermatologically acceptable carrier within which the compositions of the present invention is incorporated to enable the vitamin B<sub>3</sub> compound and preservative component, as well as other optional actives, to be delivered to the skin at an appropriate concentration.

The carrier may contain one or more dermatologically acceptable solid, semi-solid or liquid fillers, diluents, solvents, extenders and the like. The carrier may be solid, semi-solid or liquid. Preferred carriers are substantially liquid. The carrier can itself be inert or it can possess dermatological benefits of its own. Concentrations of the carrier can vary with the carrier selected and the intended concentrations of the essential and optional components.

Suitable carriers include conventional or otherwise known carriers that are dermatologically acceptable. The carrier should also be physically and chemically compatible with the essential components described herein, and should not unduly impair stability, efficacy or other use benefits associated with the compositions of the present invention. Preferred components of the compositions of this invention

should be capable of being comingled in a manner such that there is no interaction which would substantially reduce the efficacy of the composition under ordinary use situations.

The type of carrier utilized in the present invention depends on the type of product form desired for the composition. The topical compositions useful in the subject invention may be made into a wide variety of product forms such as are known in the art. These include, but are not limited to, lotions, creams, gels, sticks, sprays, ointments, pastes, mousses and cosmetics (e.g., solid, semi-solid, or liquid make-up, including foundations, eye-makeup, pigmented or non-pigmented lip treatments, e.g., lipsticks, and the like). These product forms may comprise several types of carriers including, but not limited to, solutions, aerosols, emulsions, gels, solids, and liposomes.

Preferred carriers contain a dermatologically acceptable, hydrophilic diluent. As used herein, "diluent" includes materials in which the particulate material can be dispersed, dissolved, or otherwise incorporated. Nonlimiting examples of hydrophilic diluents are water, organic hydrophilic diluents such as lower monovalent alcohols (e.g., C<sub>1</sub> - C<sub>4</sub>) and low molecular weight glycols and polyols, including propylene glycol, polyethylene glycol (e.g., Molecular Weight 200-600 g/mole), polypropylene glycol (e.g., Molecular Weight 425-2025 g/mole), glycerol, butylene glycol, 1,2,4-butanetriol, sorbitol esters, 1,2,6-hexanetriol, ethanol, isopropanol, sorbitol esters, butanediol, ether propanol, ethoxylated ethers, propoxylated ethers and combinations thereof. Water is a preferred diluent. The composition preferably comprises from about 60% to about 99.99% of the hydrophilic diluent.

Solutions according to the subject invention typically include a dermatologically acceptable hydrophilic diluent. Solutions useful in the subject invention preferably contain from about 60% to about 99.99% of the hydrophilic diluent.

Aerosols according to the subject invention can be formed by adding a propellant to a solution such as described above. Exemplary propellants include chloro-fluorinated lower molecular weight hydrocarbons. Additional propellants that are useful herein are described in Sagarin, Cosmetics Science and Technology, 2nd Edition, Vol. 2, pp. 443-465 (1972), incorporated herein by reference. Aerosols are typically applied to the skin as a spray-on product.

Preferred carriers comprise an emulsion comprising a hydrophilic phase comprising a hydrophilic component, e.g., water or other hydrophilic diluent, and a hydrophobic phase comprising a hydrophobic component, e.g., a lipid, oil or oily

material. As well known to one skilled in the art, the hydrophilic phase will be dispersed in the hydrophobic phase, or vice versa, to form respectively hydrophilic or hydrophobic dispersed and continuous phases, depending on the composition ingredients. In emulsion technology, the term "dispersed phase" is a term well-known to one skilled in the art which means that the phase exists as small particles or droplets that are suspended in and surrounded by a continuous phase. The dispersed phase is also known as the internal or discontinuous phase. The emulsion may be or comprise (e.g., in a triple or other multi-phase emulsion) an oil-in-water emulsion or a water-in-oil emulsion such as a water-in-silicone emulsion. Oil-in-water emulsions typically comprise from about 1% to about 50% of the dispersed hydrophobic phase and from about 1% to about 98% of the continuous hydrophilic phase; water-in-oil emulsions typically comprise from about 1% to about 98% of the dispersed hydrophilic phase and from about 1% to about 50% of the continuous hydrophobic phase. The emulsion may also comprise a gel network, such as described in "Application of Emulsion Stability Theories to Mobile and Semisolid Oil-in-Water Emulsions", Cosmetics and Toiletries, vol. 101, November 1986, pp. 73-92, which is incorporated by reference herein. Preferred emulsions are further described below.

The topical compositions of the subject invention, including but not limited to lotions and creams, may comprise a dermatologically acceptable emollient. Such compositions preferably contain from about 2% to about 50% of the emollient. Emollients tend to lubricate the skin, increase the smoothness and suppleness of the skin, prevent or relieve dryness of the skin, and/or protect the skin. Emollients are typically water-immiscible, oily or waxy materials. A wide variety of suitable emollients are known and may be used herein. Sagarin, Cosmetics, Science and Technology, 2nd Edition, Vol. 1, pp. 32-43 (1972), incorporated herein by reference, contains numerous examples of materials suitable as an emollient.

Lotions and creams according to the present invention generally comprise a solution carrier system and one or more emollients. Lotions typically comprise from about 1% to about 20%, preferably from about 5% to about 10%, of emollient; from about 50% to about 90%, preferably from about 60% to about 80%, water. A cream typically comprises from about 5% to about 50%, preferably from about 10% to about 20%, of emollient; and from about 45% to about 85%, preferably from about 50% to about 75%, water.

Ointments of the present invention may comprise a simple carrier base of animal or vegetable oils or semi-solid hydrocarbons (oleaginous); absorption ointment bases which absorb water to form emulsions; or water soluble carriers, e.g.,

a water soluble solution carrier. Ointments may further comprise a thickening agent, such as described in Sagarin, Cosmetics, Science and Technology, 2nd Edition, Vol. 1, pp. 72-73 (1972), incorporated herein by reference, and/or an emollient. For example, an ointment may comprise from about 2% to about 10% of an emollient; and from about 0.1% to about 2% of a thickening agent.

Compositions of this invention useful for cleansing ("cleansers") are formulated with a suitable carrier, e.g., as described above, and preferably contain one or more dermatologically acceptable surfactants in an amount which is safe and effective for cleansing. Preferred compositions contain from about 1% to about 90%, more preferably from about 5% to about 10%, of a dermatologically acceptable surfactant. The surfactant is suitably selected from anionic, cationic, nonionic, zwitterionic, amphoteric and ampholytic surfactants, as well as mixtures of these surfactants. Such surfactants are well known to those skilled in the detergency art. Nonlimiting examples of possible surfactants include isoceteth-20, sodium methyl cocoyl taurate, sodium methyl oleoyl taurate, sodium lauryl sulfate, and betaines such as described herein. See U.S. Patent No. 4,800,197, to Kowcz et al., issued January 24, 1989, which is incorporated herein by reference in its entirety, for exemplary surfactants useful herein. Examples of a broad variety of additional surfactants useful herein are described in McCutcheon's Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation, which is incorporated herein by reference in its entirety. The cleansing compositions can optionally contain, at their art-established levels, other materials which are conventionally used in cleansing compositions.

The physical form of the cleansing compositions is not critical. The compositions can be, for example, formulated as toilet bars, liquids, shampoos, bath gels, hair conditioners, hair tonics, pastes, or mousses. Toilet bars are most preferred since this is the form of cleansing agent most commonly used to wash the skin. Preferred rinse-off cleansing compositions, such as shampoos, include a delivery system adequate to deposit sufficient levels of actives on the skin and scalp. A preferred delivery system involves the use of insoluble complexes. For a more complete disclosure of such delivery systems, see U.S. Patent 4,835,148, Barford et al., issued May 30, 1989, incorporated herein by reference in its entirety.

As used herein, the term "foundation" refers to a liquid, semi-liquid, semi-solid, or solid skin cosmetic which includes, but is not limited to lotions, creams, gels, pastes, cakes, and the like. Typically the foundation is used over a large area of the skin, such as over the face, to provide a particular look. Foundations are typically used to provide an adherent base for color cosmetics such as rouge,

blusher, powder and the like, and tend to hide skin imperfections and impart a smooth, even appearance to the skin. Foundations of the present invention include a dermatologically acceptable carrier for the essential particulate material and may include conventional ingredients such as oils, colorants, pigments, emollients, fragrances, waxes, stabilizers, and the like. Exemplary carriers and such other ingredients which are suitable for use herein are described, for example, in copending patent application Serial No. 08/430,961, filed on April 28, 1995 in the names of Marcia L. Canter, Brian D. Barford, and Brian D. Hofrichter, incorporated herein by reference.

The compositions of the present invention are preferably formulated to have a pH of about 10.5 or below. The pH values of these compositions preferably range from about 2 to about 10.5, more preferably from about 3 to about 8, even more preferably from about 5 to about 8.

Preferred compositions of this invention

Preferred topical compositions of the present invention comprise an emulsion. Emulsions of the present invention may contain one or more of the following

a) Hydrophobic component

Emulsions according to the present invention contain a hydrophobic phase comprising a lipid, oil, oily or other hydrophobic component. The compositions of the present invention preferably comprise from about 1% to about 50%, preferably from about 1% to about 30%, and more preferably from about 1% to about 10% by weight of the composition of a hydrophobic component. The hydrophobic component may be derived from animals, plants, or petroleum and may be natural or synthetic (i.e., man-made). Preferred hydrophobic components are substantially water-insoluble, more preferably essentially water-insoluble. Preferred hydrophobic components are those having a melting point of about 25°C or less under about one atmosphere of pressure, and are suitable for conditioning the skin or hair.

Nonlimiting examples of suitable hydrophobic components include those selected from the group consisting of:

(1) Mineral oil, which is also known as petrolatum liquid, is a mixture of liquid hydrocarbons obtained from petroleum. See The Merck Index, Tenth Edition, Entry 7048, p. 1033 (1983) and International Cosmetic Ingredient Dictionary, Fifth Edition, vol. 1, p.415-417 (1993), which are incorporated by reference herein in their entirety.

(2) Petrolatum, which is also known as petroleum jelly, is a colloidal system of nonstraight-chain solid hydrocarbons and high-boiling liquid hydrocarbons, in which

most of the liquid hydrocarbons are held inside the micelles. See The Merck Index, Tenth Edition. Entry 7047, p. 1033 (1983); Schindler, Drug Cosmet. Ind. 89, 36-37, 76, 78-80, 82 (1961); and International Cosmetic Ingredient Dictionary, Fifth Edition, vol. 1, p. 537 (1993), which are incorporated by reference herein in their entirety.

(3) Straight and branched chain hydrocarbons having from about 7 to about 40 carbon atoms. Nonlimiting examples of these hydrocarbon materials include dodecane, isododecane, squalane, cholesterol, hydrogenated polyisobutylene, docosane (i.e. a C<sub>22</sub> hydrocarbon), hexadecane, isohexadecane (a commercially available hydrocarbon sold as Permethyl® 101A by Presperse, South Plainfield, NJ). Also useful are the C7-C40 isoparaffins, which are C7-C40 branched hydrocarbons.

(4) C1-C30 alcohol esters of C1-C30 carboxylic acids and of C2-C30 dicarboxylic acids, including straight and branched chain materials as well as aromatic derivatives (as used herein in reference to the hydrophobic component, mono- and poly- carboxylic acids include straight chain, branched chain and aryl carboxylic acids). Nonlimiting examples include diisopropyl sebacate, diisopropyl adipate, isopropyl myristate, isopropyl palmitate, methyl palmitate, myristyl propionate, 2-ethylhexyl palmitate, isodecyl neopentanoate, di-2-ethylhexyl maleate, cetyl palmitate, myristyl myristate, stearyl stearate, isopropyl stearate, methyl stearate, cetyl stearate, behenyl behenrate, dioctyl maleate, dioctyl sebacate, diisopropyl adipate, cetyl octanoate, diisopropyl dilinoleate.

(5) mono-, di- and tri- glycerides of C1-C30 carboxylic acids, e.g., caprylic/capric triglyceride, PEG-6 caprylic/capric triglyceride, PEG-8 caprylic/capric triglyceride.

(6) alkylene glycol esters of C1-C30 carboxylic acids, e.g., ethylene glycol mono- and di- esters, and propylene glycol mono- and di- esters of C1-C30 carboxylic acids e.g., ethylene glycol distearate.

(7) propoxylated and ethoxylated derivatives of the foregoing materials.

(8) C1-C30 mono- and poly- esters of sugars and related materials. These esters are derived from a sugar or polyol moiety and one or more carboxylic acid moieties. Depending on the constituent acid and sugar, these esters can be in either liquid or solid form at room temperature. Examples of liquid esters include: glucose tetraoleate, the glucose tetraesters of soybean oil fatty acids (unsaturated), the mannose tetraesters of mixed soybean oil fatty acids, the galactose tetraesters of oleic acid, the arabinose tetraesters of linoleic acid, xylose tetralinoleate, galactose pentaoleate, sorbitol tetraoleate, the sorbitol hexaesters of unsaturated soybean oil fatty acids, xylitol pentaoleate, sucrose tetraoleate, sucrose pentaoleate, sucrose



hexaoleate, sucrose heptaleate, sucrose octaoleate, and mixtures thereof. Examples of solid esters include: sorbitol hexaester in which the carboxylic acid ester moieties are palmitoleate and arachidate in a 1:2 molar ratio; the octaester of raffinose in which the carboxylic acid ester moieties are linoleate and behenate in a 1:3 molar ratio; the heptaester of maltose wherein the esterifying carboxylic acid moieties are sunflower seed oil fatty acids and lignocerate in a 3:4 molar ratio; the octaester of sucrose wherein the esterifying carboxylic acid moieties are oleate and behenate in a 2:6 molar ratio; and the octaester of sucrose wherein the esterifying carboxylic acid moieties are laurate, linoleate and behenate in a 1:3:4 molar ratio. A preferred solid material is sucrose polyester in which the degree of esterification is 7-8, and in which the fatty acid moieties are C18 mono- and/or di-unsaturated and behenic, in a molar ratio of unsaturates:behenic of 1:7 to 3:5. A particularly preferred solid sugar polyester is the octaester of sucrose in which there are about 7 behenic fatty acid moieties and about 1 oleic acid moiety in the molecule. Other materials include cottonseed oil or soybean oil fatty acid esters of sucrose. The ester materials are further described in, U.S. Patent No. 2,831,854, U.S. Patent No. 4,005,196, to Jandacek, issued January 25, 1977; U.S. Patent No. 4,005,195, to Jandacek, issued January 25, 1977; U.S. Patent No. 5,306,516, to Letton et al., issued April 26, 1994; U.S. Patent No. 5,306,515, to Letton et al., issued April 26, 1994; U.S. Patent No. 5,305,514, to Letton et al., issued April 26, 1994; U.S. Patent No. 4,797,300, to Jandacek et al., issued January 10, 1989; U.S. Patent No. 3,963,699, to Rizzi et al., issued June 15, 1976; U.S. Patent No. 4,518,772, to Volpenhein, issued May 21, 1985; and U.S. Patent No. 4,517,360, to Volpenhein, issued May 21, 1985; all of which are incorporated by reference herein in their entirety.

(9) Organopolysiloxane oils. The organopolysiloxane oil may be volatile, non-volatile, or a mixture of volatile and non-volatile silicones. The term "nonvolatile" as used in this context refers to those silicones that are liquid under ambient conditions and have a flash point (under one atmospheric pressure) of or greater than about 100°C. The term "volatile" as used in this context refers to all other silicone oils. Suitable organopolysiloxanes can be selected from a wide variety of silicones spanning a broad range of volatilities and viscosities. Nonvolatile polysiloxanes are preferred. Nonlimiting examples of suitable silicones are disclosed in U.S. Patent No. 5,069,897, to Orr, issued December 3, 1991, which is incorporated by reference herein in its entirety. Examples of suitable organopolysiloxane oils include polyalkylsiloxanes, cyclic polyalkylsiloxanes, and polyalkylarylsiloxanes.

Polyalkylsiloxanes useful in the composition herein include polyalkylsiloxanes with viscosities of from about 0.5 to about 1,000,000 centistokes at 25°C. Such polyalkylsiloxanes can be represented by the general chemical formula  $R_3SiO[R_2SiO]_xSiR_3$  wherein R is an alkyl group having from one to about 30 carbon atoms (preferably R is methyl or ethyl, more preferably methyl; also mixed alkyl groups can be used in the same molecule), and x is an integer from 0 to about 10,000, chosen to achieve the desired molecular weight which can range to over about 10,000,000. Commercially available polyalkylsiloxanes include the polydimethylsiloxanes, which are also known as dimethicones, examples of which include the Vicasil® series sold by General Electric Company and the Dow Corning® 200 series sold by Dow Corning Corporation. Specific examples of suitable polydimethylsiloxanes include Dow Corning® 200 fluid having a viscosity of 0.65 centistokes and a boiling point of 100°C, Dow Corning® 225 fluid having a viscosity of 10 centistokes and a boiling point greater than 200°C, and Dow Corning® 200 fluids having viscosities of 50, 350, and 12,500 centistokes, respectively, and boiling points greater than 200°C. Suitable dimethicones include those represented by the chemical formula  $(CH_3)_3SiO[(CH_3)_2SiO]_x[CH_3RSiO]_ySi(CH_3)_3$  wherein R is straight or branched chain alkyl having from two to about 30 carbon atoms and x and y are each integers of 1 or greater selected to achieve the desired molecular weight which can range to over about 10,000,000. Examples of these alkyl-substituted dimethicones include cetyl dimethicone and lauryl dimethicone.

Cyclic polyalkylsiloxanes suitable for use in the composition include those represented by the chemical formula  $[SiR_2-O]_n$  wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and n is an integer from about 3 to about 8, more preferably n is an integer from about 3 to about 7, and most preferably n is an integer from about 4 to about 6. When R is methyl, these materials are typically referred to as cyclomethicones. Commercially available cyclomethicones include Dow Corning® 244 fluid having a viscosity of 2.5 centistokes, and a boiling point of 172°C, which primarily contains the cyclomethicone tetramer (i.e. n=4), Dow Corning® 344 fluid having a viscosity of 2.5 centistokes and a boiling point of 178°C, which primarily contains the cyclomethicone pentamer (i.e. n=5), Dow Corning® 245 fluid having a viscosity of 4.2 centistokes and a boiling point of 205°C, which primarily contains a mixture of the cyclomethicone tetramer and pentamer (i.e. n=4 and 5), and Dow Corning® 345 fluid having a viscosity of 4.5 centistokes and a boiling point of 217°, which primarily contains a mixture of the cyclomethicone tetramer, pentamer, and hexamer (i.e. n=4, 5, and 6).

Also useful are materials such as trimethylsiloxysilicate, which is a polymeric material corresponding to the general chemical formula  $[(CH_3)_3SiO_{1/2}]_x[SiO_2]_y$ , wherein x is an integer from about 1 to about 500 and y is an integer from about 1 to about 500. A commercially available trimethylsiloxysilicate is sold as a mixture with dimethicone as Dow Corning® 593 fluid.

Dimethiconols are also suitable for use in the composition. These compounds can be represented by the chemical formulas  $R_3SiO[R_2SiO]_xSiR_2OH$  and  $HOR_2SiO[R_2SiO]_xSiR_2OH$  wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and x is an integer from 0 to about 500, chosen to achieve the desired molecular weight. Commercially available dimethiconols are typically sold as mixtures with dimethicone or cyclomethicone (e.g. Dow Corning® 1401, 1402, and 1403 fluids).

Polyalkylaryl siloxanes are also suitable for use in the composition. Polymethylphenyl siloxanes having viscosities from about 15 to about 65 centistokes at 25°C are especially useful.

Preferred for use herein are organopolysiloxanes selected from the group consisting of polyalkylsiloxanes, alkyl substituted dimethicones, cyclomethicones, trimethylsiloxysilicates, dimethiconols, polyalkylaryl siloxanes, and mixtures thereof. More preferred for use herein are polyalkylsiloxanes and cyclomethicones. Preferred among the polyalkylsiloxanes are dimethicones.

(10) Vegetable oils and hydrogenated vegetable oils. Examples of vegetable oils and hydrogenated vegetable oils include safflower oil, castor oil, coconut oil, cottonseed oil, menhaden oil, palm kernel oil, palm oil, peanut oil, soybean oil, rapeseed oil, linseed oil, rice bran oil, pine oil, sesame oil, sunflower seed oil, hydrogenated safflower oil, hydrogenated castor oil, hydrogenated coconut oil, hydrogenated cottonseed oil, hydrogenated menhaden oil, hydrogenated palm kernel oil, hydrogenated palm oil, hydrogenated peanut oil, hydrogenated soybean oil, hydrogenated rapeseed oil, hydrogenated linseed oil, hydrogenated rice bran oil, hydrogenated sesame oil, hydrogenated sunflower seed oil, and mixtures thereof.

(11) animal fats and oils, e.g., lanolin and derivatives thereof, cod liver oil.

(12) Also useful are C4-C20 alkyl ethers of polypropylene glycols, C1-C20 carboxylic acid esters of polypropylene glycols, and di-C8-C30 alkyl ethers. Nonlimiting examples of these materials include PPG-14 butyl ether, PPG-15 stearyl ether, dioctyl ether, dodecyl octyl ether, and mixtures thereof.

b) Hydrophilic component

Emulsions of the present invention also comprise a hydrophilic component, e.g., water or other hydrophilic diluent. The hydrophilic phase can thus comprise water, or a combination of water and one or more water soluble or dispersible ingredients. Hydrophilic components comprising water are preferred.

5 (c) Other components

Emulsions and other topical compositions of the present invention may comprise a variety of other ingredients such as disclosed herein. As will be understood by the skilled artisan, a given component will distribute primarily into either a hydrophilic phase or hydrophobic phase, depending on the hydrophilicity of  
10 the component in the composition.

Emulsions of the present invention preferably include one or more compounds selected from emulsifiers, surfactants, structuring agents, and thickeners.

(1) Emulsifiers/Surfactants

The emulsion may contain an emulsifier and/or surfactant, generally to help  
15 disperse and suspend the discontinuous phase within the continuous phase. A wide variety of such agents can be employed. Known or conventional emulsifiers/surfactants can be used in the composition, provided that the selected agent is chemically and physically compatible with essential components of the composition, and provides the desired dispersion characteristics. Suitable agents  
20 include non-silicone-containing emulsifiers/surfactants, silicone emulsifiers/surfactants, and mixtures thereof.


In a preferred embodiment, the composition comprises a hydrophilic emulsifier or surfactant. The compositions of the present invention preferably comprise from about 0.05% to about 5%, more preferably from about 0.05% to  
25 about 1% of at least one hydrophilic surfactant. Without intending to be limited by theory, it is believed that the hydrophilic surfactant assists in dispersing hydrophobic materials, e.g., hydrophobic structuring agents, in the hydrophilic phase. The surfactant, at a minimum, must be hydrophilic enough to disperse in the hydrophilic phase. Preferred surfactants are those having an HLB of at least about 8. The exact  
30 surfactant chosen will depend upon the pH of the composition and the other components present.

Preferred hydrophilic surfactants are selected from nonionic surfactants. Among the nonionic surfactants that are useful herein are those that can be broadly defined as condensation products of long chain alcohols, e.g. C8-30 alcohols, with  
35 sugar or starch polymers, i.e., glycosides. These compounds can be represented by the formula  $(S)_n-O-R$  wherein S is a sugar moiety such as glucose, fructose, mannose, and galactose; n is an integer of from about 1 to about 1000, and R is a

C8-30 alkyl group. Examples of long chain alcohols from which the alkyl group can be derived include decyl alcohol, cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol, oleyl alcohol, and the like. Preferred examples of these surfactants include those wherein S is a glucose moiety, R is a C8-20 alkyl group, and n is an integer of from about 1 to about 9. Commercially available examples of these surfactants include decyl polyglucoside (available as APG 325 CS from Henkel) and lauryl polyglucoside (available as APG 600 CS and 625 CS from Henkel).

Other useful nonionic surfactants include the condensation products of alkylene oxides with fatty acids (i.e. alkylene oxide esters of fatty acids). These materials have the general formula  $\text{RCO}(\text{X})_n\text{OH}$  wherein R is a C10-30 alkyl group, X is  $-\text{OCH}_2\text{CH}_2-$  (i.e. derived from ethylene glycol or oxide) or  $-\text{OCH}_2\text{CHCH}_3-$  (i.e. derived from propylene glycol or oxide), and n is an integer from about 6 to about 200. Other nonionic surfactants are the condensation products of alkylene oxides with 2 moles of fatty acids (i.e. alkylene oxide diesters of fatty acids). These materials have the general formula  $\text{RCO}(\text{X})_n\text{OOCR}$  wherein R is a C10-30 alkyl group, X is  $-\text{OCH}_2\text{CH}_2-$  (i.e. derived from ethylene glycol or oxide) or  $-\text{OCH}_2\text{CHCH}_3-$  (i.e. derived from propylene glycol or oxide), and n is an integer from about 6 to about 100. Other nonionic surfactants are the condensation products of alkylene oxides with fatty alcohols (i.e. alkylene oxide ethers of fatty alcohols). These materials have the general formula  $\text{R}(\text{X})_n\text{OR}'$  wherein R is a C10-30 alkyl group, X is  $-\text{OCH}_2\text{CH}_2-$  (i.e. derived from ethylene glycol or oxide) or  $-\text{OCH}_2\text{CHCH}_3-$  (i.e. derived from propylene glycol or oxide), and n is an integer from about 6 to about 100 and R' is H or a C10-30 alkyl group. Still other nonionic surfactants are the condensation products of alkylene oxides with both fatty acids and fatty alcohols [i.e. wherein the polyalkylene oxide portion is esterified on one end with a fatty acid and etherified (i.e. connected via an ether linkage) on the other end with a fatty alcohol]. These materials have the general formula  $\text{RCO}(\text{X})_n\text{OR}'$  wherein R and R' are C10-30 alkyl groups, X is  $-\text{OCH}_2\text{CH}_2-$  (i.e. derived from ethylene glycol or oxide) or  $-\text{OCH}_2\text{CHCH}_3-$  (derived from propylene glycol or oxide), and n is an integer from about 6 to about 100. Nonlimiting examples of these alkylene oxide derived nonionic surfactants include ceteth-6, ceteth-10, ceteth-12, cetareth-6, cetareth-10, cetareth-12, steareth-6, steareth-10, steareth-12, PEG-6 stearate, PEG-10 stearate, PEG-100 stearate, PEG-12 stearate, PEG-20 glyceryl stearate, PEG-80 glyceryl tallowate, PEG-10 glyceryl stearate, PEG-30 glyceryl cocoate, PEG-80 glyceryl cocoate, PEG-200 glyceryl tallowate, PEG-8 dilaurate, PEG-10 distearate, and mixtures thereof.

Still other useful nonionic surfactants include polyhydroxy fatty acid amide surfactants corresponding to the structural formula:



5 wherein:  $\text{R}^1$  is H,  $\text{C}_1$ - $\text{C}_4$  alkyl, 2-hydroxyethyl, 2-hydroxy-propyl, preferably  $\text{C}_1$ - $\text{C}_4$  alkyl, more preferably methyl or ethyl, most preferably methyl;  $\text{R}^2$  is  $\text{C}_5$ - $\text{C}_{31}$  alkyl or alkenyl, preferably  $\text{C}_7$ - $\text{C}_{19}$  alkyl or alkenyl, more preferably  $\text{C}_9$ - $\text{C}_{17}$  alkyl or alkenyl, most preferably  $\text{C}_{11}$ - $\text{C}_{15}$  alkyl or alkenyl; and Z is a  
10 polhydroxyhydrocarbyl moiety having a linear hydrocarbyl chain with a least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative (preferably ethoxylated or propoxylated) thereof. Z preferably is a sugar moiety selected from the group consisting of glucose, fructose, maltose, lactose, galactose, mannose, xylose, and mixtures thereof. An especially preferred surfactant corresponding to  
15 the above structure is coconut alkyl N-methyl glucoside amide (i.e., wherein the  $\text{R}^2\text{CO-}$  moiety is derived from coconut oil fatty acids). Processes for making compositions containing polyhydroxy fatty acid amides are disclosed, for example, in G.B. Patent Specification 809,060, published February 18, 1959, by Thomas Hedley & Co., Ltd.; U.S. Patent No. 2,965,576, to E. R. Wilson, issued December  
20 20, 1960; U.S. Patent No. 2,703,798, to A. M. Schwartz, issued March 8, 1955; and U.S. Patent No. 1,985,424, to Piggott, issued December 25, 1934; which are incorporated herein by reference in their entirety.

Preferred among the nonionic surfactants are those selected from the group consisting of steareth-21, cetareth-20, cetareth-12, sucrose cocoate, steareth-100,  
25 PEG-100 stearate, and mixtures thereof.

Other nonionic surfactants suitable for use herein include sugar esters and polyesters, alkoxylated sugar esters and polyesters,  $\text{C}_1$ - $\text{C}_{30}$  fatty acid esters of  $\text{C}_1$ - $\text{C}_{30}$  fatty alcohols, alkoxylated derivatives of  $\text{C}_1$ - $\text{C}_{30}$  fatty acid esters of  $\text{C}_1$ - $\text{C}_{30}$  fatty alcohols, alkoxylated ethers of  $\text{C}_1$ - $\text{C}_{30}$  fatty alcohols, polyglyceryl esters of  
30  $\text{C}_1$ - $\text{C}_{30}$  fatty acids,  $\text{C}_1$ - $\text{C}_{30}$  esters of polyols,  $\text{C}_1$ - $\text{C}_{30}$  ethers of polyols, alkyl phosphates, polyoxyalkylene fatty ether phosphates, fatty acid amides, acyl lactylates, and mixtures thereof. Nonlimiting examples of these non-silicon-containing emulsifiers include: polyethylene glycol 20 sorbitan monolaurate (Polysorbate 20), polyethylene glycol 5 soya sterol, Steareth-20, Cetareth-20, PPG-  
35 2 methyl glucose ether distearate, Ceteth-10, Polysorbate 80, cetyl phosphate.

potassium cetyl phosphate, diethanolamine cetyl phosphate, Polysorbate 60, glyceryl stearate, polyoxyethylene 20 sorbitan trioleate (Polysorbate 85), sorbitan monolaurate, polyoxyethylene 4 lauryl ether sodium stearate, polyglyceryl-4 isostearate, hexyl laurate, PPG-2 methyl glucose ether distearate, PEG-100 stearate, and mixtures thereof.

Another emulsifier useful herein are fatty acid ester blends based on a mixture of sorbitan or sorbitol fatty acid ester and sucrose fatty acid ester, the fatty acid in each instance being preferably C<sub>8</sub>-C<sub>24</sub>, more preferably C<sub>10</sub>-C<sub>20</sub>. The preferred fatty acid ester emulsifier is a blend of sorbitan or sorbitol C<sub>16</sub>-C<sub>20</sub> fatty acid ester with sucrose C<sub>10</sub>-C<sub>16</sub> fatty acid ester, especially sorbitan stearate and sucrose cocoate. This is commercially available from ICI under the trade name Arlatone 2121.

The hydrophilic surfactants useful herein can alternatively or additionally include any of a wide variety of cationic, anionic, zwitterionic, and amphoteric surfactants such as are known in the art. See, e.g., McCutcheon's, Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation; U.S. Patent No. 5,011,681 to Ciotti et al., issued April 30, 1991; U.S. Patent No. 4,421,769 to Dixon et al., issued December 20, 1983; and U.S. Patent No. 3,755,560 to Dickert et al., issued August 28, 1973; these four references are incorporated herein by reference in their entirety.

Exemplary cationic surfactants useful herein include those disclosed in U.S. Patent No. 5,151,209, to McCall et al., issued September 29, 1992; U.S. Patent No. 5,151,210, to Steuri et al., issued September 29, 1992; U.S. Patent No. 5,120,532, to Wells et al., issued June 9, 1992; U.S. Patent No. 4,387,090, to Bolich, issued June 7, 1983; U.S. Patent 3,155,591, Hilfer, issued November 3, 1964; U.S. Patent No. 3,929,678, to Laughlin et al., issued December 30, 1975; U.S. Patent No. 3,959,461, to Bailey et al., issued May 25, 1976; McCutcheon's, Detergents & Emulsifiers, (North American edition 1979) M.C. Publishing Co.; and Schwartz, et al., Surface Active Agents, Their Chemistry and Technology, New York: Interscience Publishers, 1949; all of these documents being incorporated herein by reference in their entirety. The cationic surfactants useful herein include cationic ammonium salts such as quaternary ammonium salts, and amino-amides.


A wide variety of anionic surfactants are also useful herein. See, e.g., U.S. Patent No. 3,929,678, to Laughlin et al., issued December 30, 1975, which is incorporated herein by reference in its entirety. Nonlimiting examples of anionic surfactants include the alkoyl isethionates (e.g., C<sub>12</sub> - C<sub>30</sub>), alkyl and alkyl ether sulfates and salts thereof, alkyl and alkyl ether phosphates and salts thereof, alkyl

methyl taurates (e.g., C<sub>12</sub> - C<sub>30</sub>), and soaps (e.g., alkali metal salts, e.g., sodium or potassium salts) of fatty acids.


Amphoteric and zwitterionic surfactants are also useful herein. Examples of amphoteric and zwitterionic surfactants which can be used in the compositions of the present invention are those which are broadly described as derivatives of aliphatic secondary and tertiary amines in which the aliphatic radical can be straight or branched chain and wherein one of the aliphatic substituents contains from about 8 to about 22 carbon atoms (preferably C<sub>8</sub> - C<sub>18</sub>) and one contains an anionic water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Examples are alkyl imino acetates, and iminodialkanoates and aminoalkanoates, imidazolinium and ammonium derivatives. Other suitable amphoteric and zwitterionic surfactants are those selected from the group consisting of betaines, sultaines, hydroxysultaines, alkyl sarcosinates (e.g., C<sub>12</sub> - C<sub>30</sub>), and alkanoyl sarcosinates.

Preferred emulsions of the present invention include a silicone containing emulsifier or surfactant. A wide variety of silicone emulsifiers are useful herein. These silicone emulsifiers are typically organically modified organopolysiloxanes, also known to those skilled in the art as silicone surfactants. Useful silicone emulsifiers include dimethicone copolyols. These materials are polydimethyl siloxanes which have been modified to include polyether side chains such as polyethylene oxide chains, polypropylene oxide chains, mixtures of these chains, and polyether chains containing moieties derived from both ethylene oxide and propylene oxide. Other examples include alkyl-modified dimethicone copolyols, i.e., compounds which contain C<sub>2</sub>-C<sub>30</sub> pendant side chains. Still other useful dimethicone copolyols include materials having various cationic, anionic, amphoteric, and zwitterionic pendant moieties.


The dimethicone copolyol emulsifiers useful herein can be described by the following general structure:



wherein R is C<sub>1</sub>-C<sub>30</sub> straight, branched, or cyclic alkyl and R<sup>2</sup> is selected from the group consisting of




and





wherein n is an integer from 3 to about 10; R<sup>3</sup> and R<sup>4</sup> are selected from the group consisting of H and C1-C6 straight or branched chain alkyl such that R<sup>3</sup> and R<sup>4</sup> are not simultaneously the same; and m, o, x, and y are selected such that the molecule has an overall molecular weight from about 200 to about 10,000,000, with m, o, x, and y being independently selected from integers of zero or greater such that m and o are not both simultaneously zero, and z being independently selected from integers of 1 or greater. It is recognized that positional isomers of these copolyols can be achieved. The chemical representations depicted above for the R<sup>2</sup> moieties containing the R<sup>3</sup> and R<sup>4</sup> groups are not meant to be limiting but are shown as such for convenience.

Also useful herein, although not strictly classified as dimethicone copolyols, are silicone surfactants as depicted in the structures in the previous paragraph wherein R<sup>2</sup> is:



wherein R<sup>5</sup> is a cationic, anionic, amphoteric, or zwitterionic moiety.

Nonlimiting examples of dimethicone copolyols and other silicone surfactants useful as emulsifiers herein include polydimethylsiloxane polyether copolymers with pendant polyethylene oxide sidechains, polydimethylsiloxane polyether copolymers with pendant polypropylene oxide sidechains, polydimethylsiloxane polyether copolymers with pendant mixed polyethylene oxide and polypropylene oxide sidechains, polydimethylsiloxane polyether copolymers with pendant mixed poly(ethylene)(propylene)oxide sidechains, polydimethylsiloxane polyether copolymers with pendant organobetaine sidechains, polydimethylsiloxane polyether copolymers with pendant carboxylate sidechains, polydimethylsiloxane polyether copolymers with pendant quaternary ammonium sidechains; and also further modifications of the preceding copolymers containing pendant C2-C30 straight, branched, or cyclic alkyl moieties. Examples of commercially available dimethicone copolyols useful herein sold by Dow Corning Corporation are Dow Corning® 190, 193, Q2-5220, 2501 Wax, 2-5324 fluid, and 3225C (this later material being sold as a mixture with cyclomethicone). Cetyl dimethicone copolyol is commercially available as a mixture with polyglyceryl-4 isostearate (and) hexyl laurate and is sold under the tradename ABIL® WE-09 (available from Goldschmidt). Cetyl dimethicone copolyol is also commercially available as a mixture with hexyl laurate (and) polyglyceryl-3 oleate (and) cetyl dimethicone and is sold under the tradename ABIL® WS-08 (also available from Goldschmidt). Other nonlimiting examples of dimethicone copolyols also include lauryl dimethicone copolyol, dimethicone copolyol acetate, dimethicone copolyol adipate, dimethicone

copolyolamine, dimethicone copolyol behenate, dimethicone copolyol butyl ether, dimethicone copolyol hydroxy stearate, dimethicone copolyol isostearate, dimethicone copolyol laurate, dimethicone copolyol methyl ether, dimethicone copolyol phosphate, and dimethicone copolyol stearate. See International Cosmetic  
5 Ingredient Dictionary, Fifth Edition, 1993, which is incorporated by reference herein in its entirety.

Dimethicone copolyol emulsifiers useful herein are described, for example, in U.S. Patent No. 4,960,764, to Figueroa, Jr. et al., issued October 2, 1990; European Patent No. EP 330,369, to SaNogueira, published August 30, 1989; G. H. Dahms, et al., "New Formulation Possibilities Offered by Silicone Copolyols," Cosmetics &  
10 Toiletries, vol. 110, pp. 91-100, March 1995; M. E. Carlotti et al., "Optimization of W/O-S Emulsions And Study Of The Quantitative Relationships Between Ester Structure And Emulsion Properties," J. Dispersion Science And Technology, 13(3), 315-336 (1992); P. Hameyer, "Comparative Technological Investigations of Organic and Organosilicone Emulsifiers in Cosmetic Water-in-Oil Emulsion Preparations,"  
15 HAPPI 28(4), pp. 88-128 (1991); J. Smid-Korbar et al., "Efficiency and usability of silicone surfactants in emulsions," Provisional Communication, International Journal of Cosmetic Science, 12, 135-139 (1990); and D. G. Krzysik et al., "A New Silicone Emulsifier For Water-in-Oil Systems," Drug and Cosmetic Industry, vol. 146(4) pp.  
20 28-81 (April 1990); incorporated by reference herein in their entirety.

(2) Structuring Agent

The compositions hereof, and especially the emulsions hereof, may contain a structuring agent. Structuring agents are particularly preferred in the oil-in-water emulsions of the present invention. Without being limited by theory, it is believed  
25 that the structuring agent assists in providing rheological characteristics to the composition which contribute to the stability of the composition. For example, the structuring agent tends to assist in the formation of the liquid crystalline gel network structures. The structuring agent may also function as an emulsifier or surfactant. Preferred compositions of this invention comprise from about 1% to about 20%,  
30 more preferably from about 1% to about 10%, most preferably from about 2% to about 9%, of one or more structuring agents.

Preferred structuring agents are those having an HLB of from about 1 to about 8 and having a melting point of at least about 45°C. Suitable structuring agents are those selected from the group consisting of saturated C<sub>14</sub> to C<sub>30</sub> fatty alcohols,  
35 saturated C<sub>16</sub> to C<sub>30</sub> fatty alcohols containing from about 1 to about 5 moles of ethylene oxide, saturated C<sub>16</sub> to C<sub>30</sub> diols, saturated C<sub>16</sub> to C<sub>30</sub> monoglycerol ethers, saturated C<sub>16</sub> to C<sub>30</sub> hydroxy fatty acids, C<sub>14</sub> to C<sub>30</sub> hydroxylated and

nonhydroxylated saturated fatty acids, C<sub>14</sub> to C<sub>30</sub> saturated ethoxylated fatty acids, amines and alcohols containing from about 1 to about 5 moles of ethylene oxide diols, C<sub>14</sub> to C<sub>30</sub> saturated glyceryl mono esters with a monoglyceride content of at least 40%, C<sub>14</sub> to C<sub>30</sub> saturated polyglycerol esters having from about 1 to about 3 alkyl group and from about 2 to about 3 saturated glycerol units, C<sub>14</sub> to C<sub>30</sub> glyceryl mono ethers, C<sub>14</sub> to C<sub>30</sub> sorbitan mono/diesters, C<sub>14</sub> to C<sub>30</sub> saturated ethoxylated sorbitan mono/diesters with about 1 to about 5 moles of ethylene oxide, C<sub>14</sub> to C<sub>30</sub> saturated methyl glucoside esters, C<sub>14</sub> to C<sub>30</sub> saturated sucrose mono/diesters, C<sub>14</sub> to C<sub>30</sub> saturated ethoxylated methyl glucoside esters with about 1 to about 5 moles of ethylene oxide, C<sub>14</sub> to C<sub>30</sub> saturated polyglucosides having an average of between 1 to 2 glucose units and mixtures thereof, having a melting point of at least about 45°C.

The preferred structuring agents of the present invention are selected from the group consisting of stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, stearic acid, palmitic acid, the polyethylene glycol ether of stearyl alcohol having an average of about 1 to about 5 ethylene oxide units, the polyethylene glycol ether of cetyl alcohol having an average of about 1 to about 5 ethylene oxide units, and mixtures thereof. More preferred structuring agents of the present invention are selected from the group consisting of stearyl alcohol, cetyl alcohol, behenyl alcohol, the polyethylene glycol ether of stearyl alcohol having an average of about 2 ethylene oxide units (steareth-2), the polyethylene glycol ether of cetyl alcohol having an average of about 2 ethylene oxide units, and mixtures thereof. Even more preferred structuring agents are selected from the group consisting of stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, steareth-2, and mixtures thereof.

(3) Thickening Agent (including thickeners and gelling agents)

The compositions of the present invention can also comprise a thickening agent, preferably from about 0.1% to about 5%, more preferably from about 0.1% to about 3%, and most preferably from about 0.25% to about 2%, of a thickening agent.

Nonlimiting classes of thickening agents include those selected from the group consisting of:

(i) Carboxylic Acid Polymers These polymers are crosslinked compounds containing one or more monomers derived from acrylic acid, substituted acrylic acids, and salts and esters of these acrylic acids and the substituted acrylic acids, wherein the crosslinking agent contains two or more carbon-carbon double bonds and is derived from a polyhydric alcohol. The preferred carboxylic acid polymers are of two general types. The first type of polymer is a crosslinked homopolymer of

an acrylic acid monomer or derivative thereof (e.g., wherein the acrylic acid has substituents on the two and three carbon positions independently selected from the group consisting of C<sub>1-4</sub> alkyl, -CN, -COOH, and mixtures thereof). The second type of polymer is a crosslinked copolymer having a first monomer selected from the group consisting of an acrylic acid monomer or derivative thereof (as just described in the previous sentence), a short chain alcohol (i.e., a C<sub>1-4</sub>) acrylate ester monomer or derivative thereof (e.g., wherein the acrylic acid portion of the ester has substituents on the two and three carbon positions independently selected from the group consisting of C<sub>1-4</sub> alkyl, -CN, -COOH, and mixtures thereof), and mixtures thereof; and a second monomer which is a long chain alcohol (i.e. C<sub>8-40</sub>) acrylate ester monomer or derivative thereof (e.g., wherein the acrylic acid portion of the ester has substituents on the two and three carbon positions independently selected from the group consisting of C<sub>1-4</sub> alkyl, -CN, -COOH, and mixtures thereof). Combinations of these two types of polymers are also useful herein.

In the first type of crosslinked homopolymers, the monomers are preferably selected from the group consisting of acrylic acid, methacrylic acid, ethacrylic acid, and mixtures thereof, with acrylic acid being most preferred. In the second type of crosslinked copolymers the acrylic acid monomer or derivative thereof is preferably selected from the group consisting of acrylic acid, methacrylic acid, ethacrylic acid, and mixtures thereof, with acrylic acid, methacrylic acid, and mixtures thereof being most preferred. The short chain alcohol acrylate ester monomer or derivative thereof is preferably selected from the group consisting of C<sub>1-4</sub> alcohol acrylate esters, C<sub>1-4</sub> alcohol methacrylate esters, C<sub>1-4</sub> alcohol ethacrylate esters, and mixtures thereof, with the C<sub>1-4</sub> alcohol acrylate esters, C<sub>1-4</sub> alcohol methacrylate esters, and mixtures thereof, being most preferred. The long chain alcohol acrylate ester monomer is selected from C<sub>8-40</sub> alkyl acrylate esters, with C<sub>10-30</sub> alkyl acrylate esters being preferred.

The crosslinking agent in both of these types of polymers is a polyalkenyl polyether of a polyhydric alcohol containing more than one alkenyl ether group per molecule, wherein the parent polyhydric alcohol contains at least 3 carbon atoms and at least 3 hydroxyl groups. Preferred crosslinkers are those selected from the group consisting of allyl ethers of sucrose and allyl ethers of pentaerythritol, and mixtures thereof. These polymers useful in the present invention are more fully described in U.S. Patent No. 5,087,445, to Haffey et al., issued February 11, 1992; U.S. Patent No. 4,509,949, to Huang et al., issued April 5, 1985; U.S. Patent No. 2,798,053, to Brown, issued July 2, 1957; which are both incorporated by reference herein in their entirety. See also, CTFA International Cosmetic Ingredient Dictionary, fourth

edition, 1991, pp. 12 and 80; which are also incorporated herein by reference in their entirety.

Examples of commercially available homopolymers of the first type useful herein include the carbomers, which are homopolymers of acrylic acid crosslinked with allyl ethers of sucrose or pentaerythritol. The carbomers are available as the Carbopol® 900 series from B.F. Goodrich (e.g., Carbopol® 954). Examples of commercially available copolymers of the second type useful herein include copolymers of C<sub>10-30</sub> alkyl acrylates with one or more monomers of acrylic acid, methacrylic acid, or one of their short chain (i.e. C<sub>1-4</sub> alcohol) esters, wherein the crosslinking agent is an allyl ether of sucrose or pentaerythritol. These copolymers are known as acrylates/C<sub>10-30</sub> alkyl acrylate crosspolymers and are commercially available as Carbopol® 1342, Carbopol® 1382, Pemulen TR-1, and Pemulen TR-2, from B.F. Goodrich. In other words, examples of carboxylic acid polymer thickeners useful herein are those selected from the group consisting of carbomers, acrylates/C<sub>10-C30</sub> alkyl acrylate crosspolymers, and mixtures thereof.

(ii) Crosslinked Polyacrylate Polymers The crosslinked polyacrylate polymers useful as thickeners or gelling agents include both cationic and nonionic polymers, with the cationics being generally preferred. Examples of useful crosslinked nonionic polyacrylate polymers and crosslinked cationic polyacrylate polymers are those described in U.S. Patent 5,100,660, to Hawe et al., issued March 31, 1992; U.S. Patent 4,849,484, to Heard, issued July 18, 1989; U.S. Patent 4,835,206, to Farrar et al., issued May 30, 1989; U.S. Patent 4,628,078 to Glover et al. issued December 9, 1986; U.S. Patent 4,599,379 to Flesher et al. issued July 8, 1986; and EP 228,868, to Farrar et al., published July 15, 1987; all of which are incorporated by reference herein in their entirety.

The crosslinked polyacrylate polymers are high molecular weight materials that can be characterized by the general formula: (A)<sub>l</sub>(B)<sub>m</sub>(C)<sub>n</sub> and comprise the monomer units (A)<sub>l</sub>, (B)<sub>m</sub>, and (C)<sub>n</sub>, wherein (A) is a dialkylaminoalkyl acrylate monomer or its quaternary ammonium or acid addition salt, (B) is a dialkylaminoalkyl methacrylate monomer or its quaternary ammonium or acid addition salt, (C) is a monomer that is polymerizable with (A) or (B), for example a monomer having a carbon-carbon double bond or other such polymerizable functional group, l is an integer of 0 or greater, m is an integer of 0 or greater, n is an integer of 0 or greater, but where either l or m, or both, must be 1 or greater.

The (C) monomer can be selected from any of the commonly used monomers. Nonlimiting examples of these monomers include ethylene, propylene, butylene, isobutylene, eicosene, maleic anhydride, acrylamide, methacrylamide, maleic acid,

acrolein, cyclohexene, ethyl vinyl ether, and methyl vinyl ether. In the cationic polymers of the present invention, (C) is preferably acrylamide. The alkyl portions of the (A) and (B) monomers are short chain length alkyls such as C<sub>1</sub>-C<sub>8</sub>, preferably C<sub>1</sub>-C<sub>5</sub>, more preferably C<sub>1</sub>-C<sub>3</sub>, and most preferably C<sub>1</sub>-C<sub>2</sub>. When quaternized, the polymers are preferably quaternized with short chain alkyls, i.e., C<sub>1</sub>-C<sub>8</sub>, preferably C<sub>1</sub>-C<sub>5</sub>, more preferably C<sub>1</sub>-C<sub>3</sub>, and most preferably C<sub>1</sub>-C<sub>2</sub>. The acid addition salts refer to polymers having protonated amino groups. Acid addition salts can be performed through the use of halogen (e.g. chloride), acetic, phosphoric, nitric, citric, or other acids.

These (A)<sub>l</sub>(B)<sub>m</sub>(C)<sub>n</sub> polymers also comprise a crosslinking agent, which is most typically a material containing two or more unsaturated functional groups. The crosslinking agent is reacted with the monomer units of the polymer and is incorporated into the polymer thereby forming links or covalent bonds between two or more individual polymer chains or between two or more sections of the same polymer chain. Nonlimiting examples of suitable crosslinking agents include those selected from the group consisting of methylenebisacrylamides, diallyldialkyl ammonium halides, polyalkenyl polyethers of polyhydric alcohols, allyl acrylates, vinyloxyalkylacrylates, and polyfunctional vinylidenes. Specific examples of crosslinking agents useful herein include those selected from the group consisting of methylenebisacrylamide, ethylene glycol di-(meth)acrylate, di-(meth)acrylamide, cyanomethylacrylate, vinyloxyethylacrylate, vinyloxyethylmethacrylate, allyl pentaerythritol, trimethylolpropane diallylether, allyl sucrose, butadiene, isoprene, divinyl benzene, divinyl naphthalene, ethyl vinyl ether, methyl vinyl ether, and allyl acrylate. Other crosslinkers include formaldehyde and glyoxal. Preferred for use herein as a crosslinking agent is methylenebisacrylamide.

Widely varying amounts of the crosslinking agent can be employed depending upon the properties desired in the final polymer, e.g. viscosifying effect. Without being limited by theory, it is believed that incorporation of a crosslinking agent into these cationic polymers provides a material that is a more effective viscosifying agent without negatives such as stringiness and viscosity breakdown in the presence of electrolytes. The crosslinking agent, when present, can comprise from about 1 ppm to about 1000 ppm, preferably from about 5 ppm to about 750 ppm, more preferably from about 25 ppm to about 500 ppm, even more preferably from about 100 ppm to about 500 ppm, and most preferably from about 250 ppm to about 500 ppm of the total weight of the polymer on a weight/weight basis.

The intrinsic viscosity of the crosslinked polymer, measured in one molar sodium chloride solution at 25°C, is generally above 6, preferably from about 8 to

about 14. The molecular weight (weight average) of the crosslinked polymers hereof is high, and is believed to typically be between about 1 million and about 30 million. The specific molecular weight is not critical and lower or higher weight average molecular weights can be used as long as the polymer retains its intended  
5 viscosifying effects. Preferably, a 1.0% solution of the polymer (on an actives basis) in deionized water will have a viscosity at 25°C of at least about 20,000 cP, preferably at least about 30,000 cP, when measured at 20 RPM by a Brookfield RVT (Brookfield Engineering Laboratories, Inc. Stoughton, MA, USA).

These cationic polymers can be made by polymerization of an aqueous  
10 solution containing from about 20% to about 60%, generally from about 25% to about 40%, by weight monomer, in the presence of an initiator (usually redox or thermal) until the polymerization terminates. The crosslinking agent can also be added to the solution of the monomers to be polymerized, to incorporate it into the polymer. In the polymerization reactions, the temperature generally starts between  
15 about 0° and 95°C. The polymerization can be conducted by forming a reverse phase dispersion of an aqueous phase of the monomers (and also any additional crosslinking agents) into a nonaqueous liquid, e.g. mineral oil, lanolin, isododecane, oleyl alcohol, and other volatile and nonvolatile esters, ethers, and alcohols, and the like.

20 All percentages describing the polymer in this section of the description herein are molar, unless otherwise specified. When the polymer contains (C) monomer, the molar proportion of (C) monomer, based on the total molar amount of (A), (B), and (C), can be from 0% to about 99%. The molar proportions of (A) and (B) can each be from 0% to 100%. When acrylamide, is used as the (C) monomer, it will  
25 preferably be used at a level of from about 20% to about 99%, more preferably from about 50% to about 90%.

Where monomer (A) and (B) are both present, the ratio of monomer (A) to monomer (B) in the final polymer, on a molar basis, is preferably from about 99:5 to about 15:85, more preferably from about 80:20 to about 20:80. Alternatively, in  
30 another class of polymers, the ratio is from about 5:95 to about 50:50, preferably from about 5:95 to about 25:75.

In another alternative class of polymers, the ratio (A):(B) is from about 50:50 to about 85:15. Preferably the ratio (A):(B) is about 60:40 to about 85:15, most preferably about 75:25 to about 85:15.

35 Most preferred is where monomer (A) is not present and the ratio of monomer (B):monomer (C) is from about 30:70 to about 70:30, preferably from about 40:60 to about 60:40 and most preferably from about 45:55 to about 55:45.

Cationic polymers that are useful herein that are especially preferred are those conforming to the general structure  $(A)_l(B)_m(C)_n$  wherein  $l$  is zero,  $(B)$  is methyl quaternized dimethylaminoethyl methacrylate, the ratio of  $(B):(C)$  is from about 45:55 to about 55:45, and the crosslinking agent is methylenebisacrylamide. An example of such a cationic polymer is one that is commercially available as a mineral oil dispersion (which can also include various dispersing aids such as PPG-1 trideceth-6) under the trademark Salcare® SC92 from Allied Colloids Ltd. (Norfolk, Virginia). This polymer has the proposed CTFA designation, "Polyquaternium 32 (and) Mineral Oil".

Other cationic polymers useful herein, are those not containing acrylamide or other  $(C)$  monomers, that is,  $n$  is zero. In these polymers the  $(A)$  and  $(B)$  monomer components are as described above. An especially preferred group of these non-acrylamide containing polymers is one in which  $l$  is also zero. In this instance the polymer is essentially a homopolymer of a dialkylaminoalkyl methacrylate monomer or its quaternary ammonium or acid addition salt. These dialkylaminoalkyl methacrylate polymers preferably contain a crosslinking agent as described above.

A cationic polymer, which is essentially a homopolymer, useful herein is one conforming to the general structure  $(A)_l(B)_m(C)_n$  wherein  $l$  is zero,  $(B)$  is methyl quaternized dimethylaminoethyl methacrylate,  $n$  is zero, and the crosslinking agent is methylenebisacrylamide. An example of such a homopolymer is commercially available as a mixture containing approximately 50% of the polymer, approximately 44% mineral oil, and approximately 6% PPG-1 trideceth-6 as a dispersing aid, from Allied Colloids Ltd, (Norfolk, VA) under the trademark Salcare® SC95. This polymer has recently been given the CTFA designation "Polyquaternium 37 (and) Mineral Oil (and) PPG-1 Trideceth-6".

(iii) Polyacrylamide Polymers Also useful herein are polyacrylamide polymers, especially non-ionic polyacrylamide polymers including substituted branched or unbranched polymers. These polymers can be formed from a variety of monomers including acrylamide and methacrylamide which are unsubstituted or substituted with one or two alkyl groups (preferably  $C_1$  to  $C_5$ ). Preferred are acrylate amide and methacrylate amide monomers in which the amide nitrogen is unsubstituted, or substituted with one or two  $C_1$  to  $C_5$  alkyl groups (preferably methyl, ethyl, or propyl), for example, acrylamide, methacrylamide, N-methacrylamide, N-methylmethacrylamide, N,N-dimethylmethacrylamide, N-isopropylacrylamide, N-isopropylmethacrylamide, and N,N-dimethylacrylamide. These polymers have a molecular weight greater than about 1,000,000 preferably greater than about



1.5000.000 and range up to about 30.000.000. Most preferred among these polyacrylamide polymers is the non-ionic polymer given the CTFA designation polyacrylamide and isoparaffin and laureth-7, available under the Tradename Sepigel 305 from Seppic Corporation (Fairfield, NJ).

5 Other polyacrylamide polymers useful herein include multi-block copolymers of acrylamides and substituted acrylamides with acrylic acids and substituted acrylic acids. Commercially available examples of these multi-block copolymers include Hypan SR150H, SS500V, SS500W, SSSA100H, from Lipo Chemicals, Inc., (Patterson, NJ).

10 (iv) Polysaccharides A wide variety of polysaccharides are useful herein. By "polysaccharides" are meant gelling agents containing a backbone of repeating sugar (i.e. carbohydrate) units. Nonlimiting examples of polysaccharide gelling agents include those selected from the group consisting of cellulose, carboxymethyl hydroxyethylcellulose, cellulose acetate propionate carboxylate, hydroxyethylcellulose, hydroxyethyl ethylcellulose, hydroxypropylcellulose, hydroxypropyl methylcellulose, methyl hydroxyethylcellulose, microcrystalline cellulose, sodium cellulose sulfate, and mixtures thereof. Also useful herein are the alkyl substituted celluloses. In these polymers, the hydroxy groups of the cellulose polymer is hydroxyalkylated (preferably hydroxyethylated or hydroxypropylated) to form a hydroxyalkylated cellulose which is then further modified with a C10-C30 straight chain or branched chain alkyl group through an ether linkage. Typically these polymers are ethers of C10-C30 straight or branched chain alcohols with hydroxyalkylcelluloses. Examples of alkyl groups useful herein include those selected from the group consisting of stearyl, isostearyl, lauryl, myristyl, cetyl, isocetyl, cocoyl (i.e. alkyl groups derived from the alcohols of coconut oil), palmityl, oleyl, linoleyl, linolenyl, ricinoleyl, behenyl, and mixtures thereof. Preferred among the alkyl hydroxyalkyl cellulose ethers is the material given the CTFA designation cetyl hydroxyethylcellulose, which is the ether of cetyl alcohol and hydroxyethylcellulose. This material is sold under the tradename Natrosol® CS Plus from Aqualon Corporation.

Other useful polysaccharides include scleroglucans comprising a linear chain of (1->3) linked glucose units with a (1->6) linked glucose every three units, a commercially available example of which is Clearogel™ CS11 from Michel Mercier Products Inc. (Mountainside, NJ).

35 (v) Gums Other additional thickening and gelling agents useful herein include materials which are primarily derived from natural sources. Nonlimiting examples of these gelling agent gums include materials selected from the group consisting of

acacia, agar, algin, alginic acid, ammonium alginate, amylopectin, calcium alginate, calcium carrageenan, carnitine, carrageenan, dextrin, gelatin, gellan gum, guar gum, guar hydroxypropyltrimonium chloride, hectorite, hyaluronic acid, hydrated silica, hydroxypropyl chitosan, hydroxypropyl guar, karaya gum, kelp, locust bean gum, natto gum, potassium alginate, potassium carrageenan, propylene glycol alginate, sclerotium gum, sodium carboxymethyl dextran, sodium carrageenan, tragacanth gum, xanthan gum, and mixtures thereof.

(vi) Crosslinked Vinyl Ether/Maleic Anhydride Copolymers Other additional thickening and gelling agents useful herein include crosslinked copolymers of alkyl vinyl ethers and maleic anhydride. In these copolymers the vinyl ethers are represented by the formula  $R-O-CH=CH_2$  wherein R is a C1-C6 alkyl group, preferably R is methyl. Preferred crosslinking agents are C4-C20 dienes, preferably C6 to C16 dienes, and most preferably C8 to C12 dienes. A particularly preferred copolymer is one formed from methyl vinyl ether and maleic anhydride wherein the copolymer has been crosslinked with decadiene, and wherein the polymer when diluted as a 0.5% aqueous solution at pH 7 at 25°C has a viscosity of 50,000-70,000 cps when measured using a Brookfield RTV viscometer, spindle #7 at 10 rpm. This copolymer has the CTFA designation PVM/MA decadiene crosspolymer and is commercially available as Stabileze™ 06 from International Specialty Products (Wayne NJ).

(vii) Crosslinked poly(N-vinylpyrrolidones) Crosslinked polyvinyl(N-pyrrolidones) useful herein as additional thickening and gelling agents and include those described in U.S. Patent No. 5,139,770, to Shih et al, issued August 18, 1992, and U.S. Patent No. 5,073,614, to Shih et al., issued December 17, 1991, both patents of which are incorporated by reference herein in their entirety. These gelling agents typically contain from about 0.25% to about 1% by weight of a crosslinking agent selected from the group consisting of divinyl ethers and diallyl ethers of terminal diols containing from about 2 to about 12 carbon atoms, divinyl ethers and diallyl ethers of polyethylene glycols containing from about 2 to about 600 units, dienes having from about 6 to about 20 carbon atoms, divinyl benzene, vinyl and allyl ethers of pentaerythritol, and the like. Typically, these gelling agents have a viscosity from about 25,000 cps to about 40,000 cps when measured as a 5% aqueous solution at 25 °C using a Brookfield RVT viscometer with Spindle #6 at 10 rpm. Commercially available examples of these polymers include ACP-1120, ACP-1179, and ACP-1180, available from International Specialty Products (Wayne, NJ).

Thickening agents which are suitable for use herein also include those disclosed in U.S. Patent No., 4,387,107, to Klein et al., issued June 7, 1983 and

"Encyclopedia of Polymer and Thickeners for Cosmetics." R.Y. Lochhead and W. R. Fron, eds., Cosmetics & Toiletries, vol. 108, pp. 95-135 (May 1993), which are all incorporated herein by reference in their entirety.

Preferred compositions of the present invention include a thickening agent  
5 selected from the group consisting of carboxylic acid polymers, crosslinked polyacrylate polymers, polyacrylamide polymers, and mixtures thereof, more preferably selected from the group consisting of crosslinked polyacrylate polymers, polyacrylamide polymers, and mixtures thereof.

#### Optional Components

10 The topical compositions of the present invention may comprise a wide variety of optional components, provided that such optional components are physically and chemically compatible with the essential components described herein, and do not unduly impair stability, efficacy or other use benefits associated with the compositions of the present invention. Optional components may be  
15 dispersed, dissolved or the like in the carrier of the present compositions.

Optional components include aesthetic agents and active agents. For example, the compositions may include, in addition to the essential components of the invention, absorbents (including oil absorbents such as clays and polymeric absorbents), abrasives, anticaking agents, antifoaming agents, antimicrobial agents  
20 (e.g., a compound capable of destroying microbes, preventing the development of microbes or preventing the pathogenic action of microbes and useful, for example, in controlling acne and/or preserving the topical composition), binders, biological additives, buffering agents, bulking agents, chemical additives, cosmetic biocides, denaturants, cosmetic astringents, drug astringents, external analgesics, film formers,  
25 humectants, opacifying agents, fragrances, perfumes, pigments, colorings, essential oils, skin sensates, emollients, skin soothing agents, skin healing agents, pH adjusters, plasticizers, preservatives, preservative enhancers, propellants, reducing agents, skin-conditioning agents, skin penetration enhancing agents, skin protectants, solvents, suspending agents, emulsifiers, thickening agents, solubilizing agents,  
30 polymers for aiding the film-forming properties and substantivity of the composition (such as a copolymer of eicosene and vinyl pyrrolidone, an example of which is available from GAF Chemical Corporation as Ganex® V-220), waxes, sunscreens, sunblocks, ultraviolet light absorbers or scattering agents, sunless tanning agents, antioxidants and/or radical scavengers, chelating agents, sequestrants, anti-acne  
35 agents, anti-inflammatory agents, anti-androgens, depilation agents, desquamation agents/exfoliants, organic hydroxy acids, vitamins and derivatives thereof (including water dispersible or soluble vitamins such as Vitamin C and ascorbyl phosphates).

compounds which stimulate collagen production, and natural extracts. Such other materials are known in the art. Nonexclusive examples of such materials are described in Harry's Cosmeticology, 7th Ed., Harry & Wilkinson (Hill Publishers, London 1982); in Pharmaceutical Dosage Forms- Disperse Systems; Lieberman, Rieger & Banker, Vols. 1 (1988) & 2 (1989); Marcel Decker, Inc.; in The Chemistry and Manufacture of Cosmetics, 2nd. Ed., deNavarre (Van Nostrand 1962-1965); and in The Handbook of Cosmetic Science and Technology, 1st Ed., Knowlton & Pearce (Elsevier 1993), can also be used in the present invention.

In a preferred embodiment, the composition also includes an active useful for chronically regulating skin condition. Such materials are those which manifest skin appearance benefits following chronic application of the composition containing such materials. Materials having this effect include, but are not limited to, retinoids.

A. Retinoids

In a preferred embodiment, the compositions of the present invention contain a retinoid. The retinoid enhances the skin appearance benefits of the present invention, especially in regulating skin condition, including regulating signs of skin aging, more especially wrinkles, lines, and pores.

As used herein, "retinoid" includes all natural and/or synthetic analogs of Vitamin A or retinol-like compounds which possess the biological activity of Vitamin A in the skin as well as the geometric isomers and stereoisomers of these compounds. The retinoid is preferably retinol, retinol esters (e.g., C<sub>2</sub> - C<sub>22</sub> alkyl esters of retinol, including retinyl palmitate, retinyl acetate, retinyl propionate), retinal, and/or retinoic acid (including all-trans retinoic acid and/or 13-cis-retinoic acid), more preferably retinoids other than retinoic acid. These compounds are well known in the art and are commercially available from a number of sources, e.g., Sigma Chemical Company (St. Louis, MO), and Boehringer Mannheim (Indianapolis, IN). Other retinoids which are useful herein are described in U.S. Patent Nos. 4,677,120, issued Jun. 30, 1987 to Parish et al.; 4,885,311, issued Dec. 5, 1989 to Parish et al.; 5,049,584, issued Sep. 17, 1991 to Purcell et al.; 5,124,356, issued Jun. 23, 1992 to Purcell et al.; and Reissue 34,075, issued Sep. 22, 1992 to Purcell et al.. Other suitable retinoids are tocopheryl-retinoate [tocopherol ester of retinoic acid (trans- or cis-), adapalene {6-[3-(1-adamantyl)-4-methoxyphenyl]-2-naphthoic acid}, and tazarotene (ethyl 6-[2-(4,4-dimethylthiochroman-6-yl)-ethynyl]nicotinate). One or more retinoids may be used herein. Preferred retinoids are retinol, retinyl palmitate, retinyl acetate, retinyl propionate, retinal and combinations thereof. More preferred are retinol and retinyl palmitate.

The retinoid may be included as the substantially pure material, or as an extract obtained by suitable physical and/or chemical isolation from natural (e.g., plant) sources. The retinoid is preferably substantially pure, more preferably essentially pure.

5       The compositions of this invention may contain a safe and effective amount of the retinoid, such that the resultant composition is safe and effective for regulating skin condition, preferably for regulating visible and/or tactile discontinuities in skin, more preferably for regulating signs of skin aging, even more preferably for regulating visible and/or tactile discontinuities in skin texture associated with skin  
10   aging. The compositions preferably contain from or about 0.005% to or about 2%, more preferably 0.01% to or about 2%, retinoid. Retinol is most preferably used in an amount of from or about 0.01% to or about 0.15%; retinol esters are most preferably used in an amount of from or about 0.01% to or about 2% (e.g., about 1%); retinoic acids are most preferably used in an amount of from or about 0.01% to  
15   or about 0.25%; tocopheryl-retinoate, adapalene, and tazarotene are most preferably used in an amount of from or about 0.01% to or about 2%.

In a preferred embodiment, the composition contains both a retinoid and a Vitamin B<sub>3</sub> compound. The retinoid is preferably used in the above amounts, and the vitamin B<sub>3</sub> compound is preferably used in an amount of from or about 0.1% to  
20   or about 10%, more preferably from or about 2% to or about 5%.

#### B. Anti-Inflammatory Agents

A safe and effective amount of an anti-inflammatory agent may be added to the compositions of the subject invention, preferably from about 0.1% to about 10%, more preferably from about 0.5% to about 5%, of the composition. The anti-  
25   inflammatory agent enhances the skin appearance benefits of the present invention, e.g., such agents contribute to a more uniform and acceptable skin tone or color. The exact amount of anti-inflammatory agent to be used in the compositions will depend on the particular anti-inflammatory agent utilized since such agents vary widely in potency.

30       Steroidal anti-inflammatory agents, including but not limited to, corticosteroids such as hydrocortisone, hydroxyltriamcinolone, alpha-methyl dexamethasone, dexamethasone-phosphate, beclomethasone dipropionates, clobetasol valerate, desonide, desoxymethasone, desoxycorticosterone acetate, dexamethasone, dichlorisone, diflorasone diacetate, diflucortolone valerate,  
35   fluadrenolone, flucorolone acetonide, fludrocortisone, flumethasone pivalate, fluosinolone acetonide, fluocinonide, flucortine butylesters, fluocortolone, fluprednidene (fluprednylidene) acetate, flurandrenolone, halcinonide,

hydrocortisone acetate, hydrocortisone butyrate, methylprednisolone, triamcinolone acetonide, cortisone, cortodoxone, flucetonide, fludrocortisone, difluorosone diacetate, fluradrenolone, fludrocortisone, difluorosone diacetate, fluradrenolone acetonide, medrysone, amcinafel, amcinafide, betamethasone and the balance of its esters, chlorprednisone, chlorprednisone acetate, clocortelone, clescinolone, dichlorisone, diflurprednate, flucoronide, flunisolide, fluoromethalone, fluperolone, fluprednisolone, hydrocortisone valerate, hydrocortisone cyclopentylpropionate, hydrocortamate, meprednisone, paramethasone, prednisolone, prednisone, beclomethasone dipropionate, triamcinolone, and mixtures thereof may be used.

10 The preferred steroidal anti-inflammatory for use is hydrocortisone.

A second class of anti-inflammatory agents which is useful in the compositions includes the nonsteroidal anti-inflammatory agents. The variety of compounds encompassed by this group are well-known to those skilled in the art. For detailed disclosure of the chemical structure, synthesis, side effects, etc. of non-steroidal anti-inflammatory agents, reference may be had to standard texts, including

15 Anti-inflammatory and Anti-Rheumatic Drugs, K. D. Rainsford, Vol. I-III, CRC Press, Boca Raton, (1985), and Anti-inflammatory Agents, Chemistry and Pharmacology, I. R. A. Scherrer, et al., Academic Press, New York (1974), each incorporated herein by reference.

20 Specific non-steroidal anti-inflammatory agents useful in the composition invention include, but are not limited to:

- 1) the oxicams, such as piroxicam, isoxicam, tenoxicam, sudoxicam, and CP-14,304;
- 2) the salicylates, such as aspirin, disalcid, benorylate, trilisate, safapryn,
- 25 solprin, diflunisal, and fendosal;
- 3) the acetic acid derivatives, such as diclofenac, fenclofenac, indomethacin, sulindac, tolmetin, isoxepac, furofenac, tiopinac, zidometacin, acematacin, fentiazac, zomepirac, clindanac, oxepinac, felbinac, and ketorolac;
- 4) the fenamates, such as mefenamic, meclofenamic, flufenamic,
- 30 niflumic, and tolfenamic acids;
- 5) the propionic acid derivatives, such as ibuprofen, naproxen, benoxaprofen, flurbiprofen, ketoprofen, fenoprofen, fenbufen, indoprofen, piroprofen, carprofen, oxaprozin, pranoprofen, miroprofen, tioxaprofen, suprofen, alminoprofen, and tiaprofenic; and
- 35 6) the pyrazoles, such as phenylbutazone, oxyphenbutazone, feprazone, azapropazone, and trimethazone.

Mixtures of these non-steroidal anti-inflammatory agents may also be employed, as well as the dermatologically acceptable salts and esters of these agents. For example, etofenamate, a flufenamic acid derivative, is particularly useful for topical application. Of the nonsteroidal anti-inflammatory agents, ibuprofen, naproxen, flufenamic acid, etofenamate, aspirin, mefenamic acid, meclofenamic acid, piroxicam and felbinac are preferred; ibuprofen, naproxen, etofenamate, aspirin and flufenamic acid are most preferred.

Finally, so-called "natural" anti-inflammatory agents are useful in methods of the subject invention. Such agents may suitably be obtained as an extract by suitable physical and/or chemical isolation from natural sources (e.g., plants, fungi, by-products of microorganisms). For example, candelilla wax, alpha bisabolol, aloe vera, Manjistha (extracted from plants in the genus Rubia, particularly Rubia Cordifolia), and Guggal (extracted from plants in the genus Commiphora, particularly Commiphora Mukul), kola extract, chamomile, and sea whip extract, may be used.

Additional anti-inflammatory agents useful herein include compounds of the Licorice (the plant genus/species Glycyrrhiza glabra) family, including glycyrrhetic acid, glycyrrhizic acid, and derivatives thereof (e.g., salts and esters). Suitable salts of the foregoing compounds include metal and ammonium salts. Suitable esters include C<sub>2</sub> - C<sub>24</sub> saturated or unsaturated esters of the acids, preferably C<sub>10</sub> - C<sub>24</sub>, more preferably C<sub>16</sub> - C<sub>24</sub>. Specific examples of the foregoing include oil soluble licorice extract, the glycyrrhizic and glycyrrhetic acids themselves, monoammonium glycyrrhizinate, monopotassium glycyrrhizinate, dipotassium glycyrrhizinate, 1-beta-glycyrrhetic acid, stearyl glycyrrhetinate, and 3-stearyloxy-glycyrrhetinic acid, and disodium 3-succinyloxy-beta-glycyrrhetinate. Stearyl glycyrrhetinate is preferred.

#### C. Sunscreens and Sunblocks

Exposure to ultraviolet light can result in excessive scaling and texture changes of the stratum corneum. Therefore, the compositions of the subject invention preferably contain a sunscreen or sunblock. Suitable sunscreens or sunblocks may be organic or inorganic.

A wide variety of conventional suncreening agents are suitable for use herein. Sagarin, et al., at Chapter VIII, pages 189 et seq., of Cosmetics Science and Technology (1972), discloses numerous suitable agents, and is incorporated herein by reference. Specific suitable suncreening agents include, for example: p-aminobenzoic acid, its salts and its derivatives (ethyl, isobutyl, glyceryl esters; p-

dimethylaminobenzoic acid); anthranilates (i.e., o-amino-benzoates; methyl, menthyl, phenyl, benzyl, phenylethyl, linalyl, terpinyl, and cyclohexenyl esters); salicylates (amyl, phenyl, octyl, benzyl, menthyl, glyceryl, and di-pro-pyleneglycol esters); cinnamic acid derivatives (menthyl and benzyl esters, a-phenyl cinnamionitrile; butyl cinnamoyl pyruvate); dihydroxycinnamic acid derivatives (umbelliferone, methylumbelliferone, methylaceto-umbelliferone); trihydroxycinnamic acid derivatives (esculetin, methylesculetin, daphnetin, and the glucosides, esculin and daphnin); hydrocarbons (diphenylbutadiene, stilbene); dibenzalacetone and benzalacetophenone; naphtholsulfonates (sodium salts of 2-naphthol-3,6-disulfonic and of 2-naphthol-6,8-disulfonic acids); di-hydroxynaphthoic acid and its salts; o- and p-hydroxybiphenyldisulfonates; coumarin derivatives (7-hydroxy, 7-methyl, 3-phenyl); diazoles (2-acetyl-3-bromoindazole, phenyl benzoxazole, methyl naphthoxazole, various aryl benzothiazoles); quinine salts (bisulfate, sulfate, chloride, oleate, and tannate); quinoline derivatives (8-hydroxyquinoline salts, 2-phenylquinoline); hydroxy- or methoxy-substituted benzophenones; uric and violuric acids; tannic acid and its derivatives (e.g., hexaethylether); (butyl carbotol) (6-propyl piperonyl) ether; hydroquinone; benzophenones (oxybenzene, sulisobenzene, dioxybenzene, benzoescorcinol, 2,2',4,4'-tetrahydroxybenzophenone, 2,2'-dihydroxy-4,4'-dimethoxybenzophenone, octabenzene; 4-isopropylidibenzoylmethane; butylmethoxydibenzoylmethane; etocrylene; octocrylene; [3-(4'-methylbenzylidene bornan-2-one) and 4-isopropyl-dibenzoylmethane.

Of these, 2-ethylhexyl-p-methoxycinnamate (commercially available as PARSOL MCX), 4,4'-t-butyl methoxydibenzoyl-methane (commercially available as PARSOL 1789), 2-hydroxy-4-methoxybenzophenone, octyldimethyl-p-aminobenzoic acid, digalloyltriolate, 2,2-dihydroxy-4-methoxybenzophenone, ethyl-4-(bis(hydroxy-propyl))aminobenzoate, 2-ethylhexyl-2-cyano-3,3-diphenylacrylate, 2-ethylhexyl-salicylate, glyceryl-p-aminobenzoate, 3,3,5-trimethylcyclohexylsalicylate, methylanthranilate, p-dimethyl-aminobenzoic acid or aminobenzoate, 2-ethylhexyl-p-dimethyl-amino-benzoate, 2-phenylbenzimidazole-5-sulfonic acid, 2-(p-dimethylaminophenyl)-5-sulfonicbenzoxazoic acid, octocrylene and mixtures of these compounds, are preferred.

More preferred organic sunscreens useful in the compositions useful in the subject invention are 2-ethylhexyl-p-methoxycinnamate, butylmethoxydibenzoylmethane, 2-hydroxy-4-methoxybenzo-phenone, 2-phenylbenzimidazole-5-sulfonic acid, octyldimethyl-p-aminobenzoic acid, octocrylene and mixtures thereof.



Also particularly useful in the compositions are sunscreens such as those disclosed in U.S. Patent No. 4,937,370 issued to Sabatelli on June 26, 1990, and U.S. Patent No. 4,999,186 issued to Sabatelli & Spirnak on March 12, 1991, both of which are incorporated herein by reference. The suncreening agents disclosed  
5 therein have, in a single molecule, two distinct chromophore moieties which exhibit different ultra-violet radiation absorption spectra. One of the chromophore moieties absorbs predominantly in the UVB radiation range and the other absorbs strongly in the UVA radiation range.

Preferred members of this class of suncreening agents are 4-N,N-(2-ethylhexyl)methyl-aminobenzoic acid ester of 2,4-dihydroxybenzophenone; N,N-di-  
10 (2-ethylhexyl)-4-aminobenzoic acid ester with 4-hydroxydibenzoylmethane; 4-N,N-(2-ethylhexyl)methyl-aminobenzoic acid ester with 4-hydroxydibenzoylmethane; 4-N,N-(2-ethylhexyl)methyl-aminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy)benzophenone; 4-N,N-(2-ethylhexyl)-methylaminobenzoic acid ester  
15 of 4-(2-hydroxyethoxy)dibenzoylmethane; N,N-di-(2-ethylhexyl)-4-aminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy)benzophenone; and N,N-di-(2-ethylhexyl)-4-aminobenzoic acid ester of 4-(2-hydroxyethoxy)dibenzoylmethane and mixtures thereof.

Especially preferred suncreens or sunblocks include  
20 butylmethoxydibenzoylmethane, 2-ethylhexyl-p-methoxycinnamate, phenyl benzimidazole sulfonic acid, and octocrylene.

A safe and effective amount of the sunscreen or sunblock is used, typically from about 1% to about 20%, more typically from about 2% to about 10%. Exact amounts will vary depending upon the sunscreen chosen and the desired Sun  
25 Protection Factor (SPF).

An agent may also be added to any of the compositions useful in the subject invention to improve the skin substantivity of those compositions, particularly to enhance their resistance to being washed off by water, or rubbed off. A preferred agent which will provide this benefit is a copolymer of ethylene and acrylic acid.  
30 Compositions comprising this copolymer are disclosed in U.S. Patent 4,663,157, Brock, issued May 5, 1987, which is incorporated herein by reference.

#### D. Anti-Oxidants/Radical Scavengers

Preferred compositions of the subject invention include an anti-oxidant/radical scavenger. The anti-oxidant/radical scavenger is especially useful  
35 for providing protection against UV radiation which can cause increased scaling or texture changes in the stratum corneum and against other environmental agents which can cause skin damage.

A safe and effective amount of an anti-oxidant/radical scavenger may be added to the compositions of the subject invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition.

Anti-oxidants/radical scavengers such as ascorbic acid (vitamin C) and its salts, ascorbyl esters of fatty acids, ascorbic acid derivatives (e.g., magnesium ascorbyl phosphate), tocopherol (vitamin E), tocopherol sorbate, tocopherol acetate, other esters of tocopherol, butylated hydroxy benzoic acids and their salts, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (commercially available under the tradename Trolox<sup>R</sup>), gallic acid and its alkyl esters, especially propyl gallate, uric acid and its salts and alkyl esters, sorbic acid and its salts, amines (e.g., N,N-diethylhydroxylamine, amino-guanidine), sulfhydryl compounds (e.g., glutathione), dihydroxy fumaric acid and its salts, lysine pidolate, arginine pidolate, nordihydroguaiaretic acid, bioflavonoids, lysine, methionine, proline, superoxide dismutase, silymarin, tea extracts, grape skin/seed extracts, melanin, and rosemary extracts may be used. Preferred anti-oxidants/radical scavengers are selected from tocopherol sorbate and other esters of tocopherol, more preferably tocopherol sorbate. For example, the use of tocopherol sorbate in topical compositions and applicable to the present invention is described in U.S. Patent No. 4,847,071, issued on July 11, 1989 to Donald L. Bissett, Rodney D. Bush and Ranjit Chatterjee, incorporated herein by reference.

#### E. Chelators

As used herein, "chelating agent" means an active agent capable of removing a metal ion from a system by forming a complex so that the metal ion cannot readily participate in or catalyze chemical reactions. The inclusion of a chelating agent is especially useful for providing protection against UV radiation which can contribute to excessive scaling or skin texture changes and against other environmental agents which can cause skin damage.

A safe and effective amount of a chelating agent may be added to the compositions of the subject invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition. Exemplary chelators that are useful herein are disclosed in U.S. Patent No. 5,487,884, issued 1/30/96 to Bissett et al.; International Publication No. 91/16035, Bush et al., published 10/31/95; and International Publication No. 91/16034, Bush et al., published 10/31/95; all incorporated herein by reference. Preferred chelators useful in compositions of the subject invention are furildioxime and derivatives thereof.

#### F. Organic Hydroxy Acids

Compositions of the present invention may comprise an organic hydroxy acid. Suitable hydroxy acids include C<sub>1</sub> - C<sub>18</sub> hydroxy acids, preferably C<sub>8</sub> or below. The hydroxy acids can be substituted or unsubstituted, straight chain, branched chain or cyclic (preferably straight chain), and saturated or unsaturated (mono- or poly- unsaturated) (preferably saturated). Non-limiting examples of suitable hydroxy acids include salicylic acid, glycolic acid, lactic acid, 5 octanoyl salicylic acid, hydroxyoctanoic acid, hydroxycaprylic acid, and lanolin fatty acids. Preferred concentrations of the organic hydroxy acid range from about 0.1% to about 10%, more preferably from about 0.2% to about 5%, also preferably from about 0.5% to about 2%. Salicylic acid is preferred. The organic hydroxy acids enhance the skin appearance benefits of the present invention. For example, the organic hydroxy acids tend to improve the texture of the skin.

#### G. Desquamation Agents/Exfoliants

A safe and effective amount of a desquamation agent may be added to the compositions of the subject invention, more preferably from about 0.1% to about 10%, even more preferably from about 0.2% to about 5%, also preferably from about 0.5% to about 4% of the composition. Desquamation agents enhance the skin appearance benefits of the present invention. For example, the desquamation agents tend to improve the texture of the skin (e.g., smoothness). A variety of desquamation agents are known in the art and are suitable for use herein, including but not limited to the organic hydroxy agents described above. One desquamation system that is suitable for use herein comprises sulfhydryl compounds and zwitterionic surfactants and is described in copending application Serial No. 08/480,632, filed on June 7, 1995 in the name of Donald L. Bissett, corresponding to PCT Application No. U.S. 95/08136, filed 6/29/95, each incorporated herein by reference. Another desquamation system that is suitable for use herein comprises salicylic acid and zwitterionic surfactants and is described in copending patent application Serial No. 08/554,944, filed on November 13, 1995 as a continuation of Serial No. 08/209,401, filed on March 9, 1994 in the name of Bissett, corresponding to PCT Application No. 94/12745, filed 11/4/94, published 5/18/95, each incorporated herein by reference. Zwitterionic surfactants such as described in these applications are also useful as desquamatory agents herein, with cetyl betaine being particularly preferred.

#### H. Skin Lightening Agents

The compositions of the present invention may comprise a skin lightening agent. When used, the compositions preferably comprise from about 0.1% to about

10%, more preferably from about 0.2% to about 5%, also preferably from about 0.5% to about 2%, of a skin lightening agent. Suitable skin lightening agents include those known in the art, including kojic acid, arbutin, ascorbic acid and derivatives thereof, e.g., magnesium ascorbyl phosphate. Skin lightening agents  
5 suitable for use herein also include those described in copending patent application Serial No. 08/479,935, filed on June 7, 1995 in the name of Hillebrand, corresponding to PCT Application No. U.S. 95/07432, filed 6/12/95; and copending patent application Serial No. 08/390,152, filed on February 24, 1995 in the names of Kalla L. Kvalnes, Mitchell A. DeLong, Barton J. Bradbury, Curtis B. Motley, and  
10 John D. Carter, corresponding to PCT Application No. U.S. 95/02809, filed 3/1/95, published 9/8/95; all incorporated herein by reference.

I. Skin Conditioners

Preferred compositions of the invention comprise an optional skin conditioning component. The skin conditioning component is preferably selected  
15 from the group consisting of emollients, humectants, moisturizers and mixtures thereof. The skin conditioning component is preferably present at a level of at least about 0.1%, more preferably from about 1% to about 99%, even more preferably from about 1% to about 50%, still more preferably from about 2% to about 30% and most preferably from about 5% to about 25% (e.g., about 5% to about 10% or 15%).

20 A variety of emollients may be employed. These emollients may be selected from one or more of the following classes: Triglyceride esters which include, but are not limited to, vegetable and animal fats and oils such as castor oil, cocoa butter, safflower oil, cottonseed oil, corn oil, olive oil, cod liver oil, almond oil, avocado oil, palm oil, sesame oil, squalene, kukui oil and soybean oil; Acetoglyceride esters,  
25 such as acetylated monoglycerides; Ethoxylated glycerides, such as ethoxylated glyceryl monostearate; Alkyl esters of fatty acids having 10 to 20 carbon atoms which include, but are not limited to, methyl, isopropyl, and butyl esters of fatty acids such as hexyl laurate, isohexyl laurate, isohexyl palmitate, isopropyl palmitate, methyl palmitate, decyloleate, isodecyl oleate, hexadecyl stearate decyl stearate,  
30 isopropyl isostearate, methyl isostearate, diisopropyl adipate, diisohexyl adipate, dihexyldecyl adipate, diisopropyl sebacate, lauryl lactate, myristyl lactate, and cetyl lactate; Alkenyl esters of fatty acids having 10 to 20 carbon atoms such as oleyl myristate, oleyl stearate, and oleyl oleate; Fatty acids having 10 to 20 carbon atoms such as pelargonic, lauric, myristic, palmitic, stearic, isostearic, hydroxystearic,  
35 oleic, linoleic, ricinoleic, arachidic, behenic, and erucic acids; Fatty alcohols having 10 to 20 carbon atoms such as lauryl, myristyl, cetyl, hexadecyl, stearyl, isostearyl, hydroxystearyl, oleyl, ricinoleyl, behenyl, erucyl, and 2-octyl dodecanyl alcohols;

Lanolin and lanolin derivatives such as lanolin, lanolin oil, lanolin wax, lanolin alcohols, lanolin fatty acids, isopropyl lanolate, ethoxylated cholesterol, propoxylated lanolin alcohols, acetylated lanolin alcohols, lanolin alcohols linoleate, lanolin alcohols ricinoleate, acetate of lanolin alcohols ricinoleate, acetate of  
5 ethoxylated alcohols-esters, hydrogenolysis of lanolin, ethoxylated hydrogenated lanolin, and liquid and semisolid lanolin absorption bases; Polyhydric alcohol esters such as ethylene glycol mono and di-fatty acid esters, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol (200-6000) mono- and di-fatty acid esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol 2000  
10 monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol polyfatty esters, ethoxylated glyceryl monostearate, 1,2-butylene glycol monostearate, 1,2-butylene glycol distearate, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters; Wax esters such as beeswax, spermaceti, myristyl myristate, stearyl stearate;  
15 Beeswax derivatives such as polyoxyethylene sorbitol beeswax which are reaction products of beeswax with ethoxylated sorbitol of varying ethylene oxide content, forming a mixture of ether esters; Vegetable waxes including, but not limited to, carnauba and candelilla waxes; Phospholipids such as lecithin and derivatives; Sterols including, but not limited to, cholesterol and cholesterol fatty acid esters; and  
20 Amides such as fatty acid amides, ethoxylated fatty acid amides, and solid fatty acid alkanolamides.

Additional types of conditioning compounds are humectants of the polyhydric alcohol-type. Typical polyhydric alcohols include polyalkylene glycols and more preferably alkylene polyols and their derivatives, including propylene  
25 glycol, dipropylene glycol, polypropylene glycol, polyethylene glycol and derivatives thereof, sorbitol, hydroxypropyl sorbitol, erythritol, threitol, pentaerythritol, xylitol, glucitol, mannitol, hexylene glycol, butylene glycol (e.g., 1,3-butylene glycol), hexane triol (e.g., 1,2,6-hexanetriol), glycerol, ethoxylated glycerol, propoxylated glycerol, sodium 2-pyrrolidone-5-carboxylate, soluble  
30 collagen, dibutyl phthalate, gelatin and mixtures thereof.

Also useful herein are guanidine; glycolic acid and glycolate salts (e.g. ammonium and quaternary alkyl ammonium); lactic acid and lactate salts (e.g. ammonium and quaternary alkyl ammonium); aloe vera in any of its variety of forms (e.g., aloe vera gel); sugar and starch derivatives (e.g., alkoxyated glucose);  
35 hyaluronic acid and derivatives thereof (e.g., salt derivatives such as sodium hyaluronate); lactamide monoethanolamine; acetamide monoethanolamine; urea; panthenol; sugars; starches; silicone fluids; silicone gums; and mixtures thereof.

Also useful are the propoxylated glycerols described in U.S. Patent No. 4,976,953, which is description is incorporated herein by reference. Other useful conditioning compounds include the various C<sub>1</sub>-C<sub>30</sub> monoesters and polyesters of sugars and related materials such as described herein in reference to the hydrophobic component.

The above listed compounds may be incorporated singly or in combination.

#### Preparation of Compositions

The compositions of the present invention are generally prepared by conventional methods such as are known in the art of making topical compositions. Such methods typically involve mixing of the ingredients in one or more steps to a relatively uniform state, with or without heating, cooling, application of vacuum, and the like.

#### Methods for Regulating Skin Condition

The compositions of the present invention are useful for regulating mammalian skin condition (especially human skin, more especially human facial skin), including visible and/or tactile discontinuities in skin, signs of skin aging, and visible and/or tactile discontinuities in skin associated with skin aging (including fine lines, wrinkles, large pores, surface roughness and other texture discontinuities associated with aged skin). Such regulation includes prophylactic and therapeutic regulation.

Regulating skin condition involves topically applying to the skin a safe and effective amount of a composition of the present invention. The amount of the composition which is applied, the frequency of application and the period of use will vary widely depending upon the level of vitamin B<sub>3</sub> compound and/or other components of a given composition and the level of regulation desired, e.g., in light of the level of skin aging present in the subject and the rate of further skin aging.

In a preferred embodiment, the composition is chronically applied to the skin. By "chronic topical application" is meant continued topical application of the composition over an extended period during the subject's lifetime, preferably for a period of at least about one week, more preferably for a period of at least about one month, even more preferably for at least about three months, even more preferably for at least about six months, and more preferably still for at least about one year. While benefits are obtainable after various maximum periods of use (e.g., five, ten or twenty years), it is preferred that chronic application continue throughout the subject's lifetime. Typically applications would be on the order of about once per day over such extended periods, however application rates can vary from about once per week up to about three times per day or more.

A wide range of quantities of the compositions of the present invention can be employed to provide a skin appearance and/or feel benefit. Quantities of the present compositions which are typically applied per application are, in mg composition/cm<sup>2</sup> skin, from about 0.1 mg/cm<sup>2</sup> to about 10 mg/cm<sup>2</sup>. A particularly useful application amount is about 2 mg/cm<sup>2</sup>.

Regulating skin condition is preferably practiced by applying a composition in the form of a skin lotion, cream, cosmetic, or the like which is intended to be left on the skin for some esthetic, prophylactic, therapeutic or other benefit (i.e., a "leave-on" composition). After applying the composition to the skin, it is preferably left on the skin for a period of at least about 15 minutes, more preferably at least about 30 minutes, even more preferably at least about 1 hour, most preferably for at least several hours, e.g., up to about 12 hours.

### EXAMPLES

The following examples further describe and demonstrate embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention.

#### Example 1

A stable topical composition is prepared from the following ingredients using conventional formulating techniques.

Phase A	Ingredient	Weight Percent
	DI water	14.699
	glycerin	10.000
Phase B		
	Carbopol 954	0.400
	disodium EDTA	0.100
Phase C		
	isopropyl palmitate	0.400
	cetyl alcohol	2.300
	cyclomethicone/dimethicone copolyol	1.900
	stearyl alcohol	1.500
	dimethicone (200 cts)	0.600
	PEG 100 Stearate	0.300
	stearic acid	0.300
	cetyl palmitate	2.500
Phase D		

	DI water	3.000
	sodium hydroxide	0.200
Phase E		
	DI water	8.000
	Niacinamide	2.000
Phase F		
	Sodium hydroxy glycinate	0.100
	Isopropynyl butylcarbamide	0.100
	DI water	0.300
	butylene glycol	0.300
Phase G		
	isopropyl palmitate	1.000
	retinol	0.050
	BHT	0.001
	Polysorbate 20	0.050

First, Phase A ingredients are combined and sparged using nitrogen for approximately 15 minutes. Phase B ingredients are then dispersed into Phase A until uniform using propeller type mixing and heated the mixture to about 75°C. In a separate vessel, Phase C ingredients are combined and heated to about 75°C. The mixture of phases A and B are then blanketed with a slow, steady stream of nitrogen. Next the Phase C ingredients are homogenized into the mixture of phases A and B using any rotor/stator type of homogenizer for approximately 15 minutes. After the 15 minutes, the mixing is switched to low rpm sweep mixing. Next, phase D ingredients are combined and added to the mixture of phases A,B, and C. Once phase D is mixed and the batch mixture is homogeneous, the entire batch mixture is cooled. When the batch is cooled to about 50°C, phase E ingredients are added and homogenized. When the batch is cooled to about 40°C, phase F ingredients are added to the batch mixture. Lastly, when the batch mixture is cooled to about 30°C, the phase G ingredients are combined to the batch mixture. Mixing is continued until the batch mixture is uniform.

The resulting composition is useful for application to the skin for delivering the retinol and to treat and improve the appearance of the skin.



Example 2

A stable topical composition is prepared from the following ingredients using conventional formulating techniques.

5

Phase A	Ingredient	Weight Percent
	DI Water	
	disodium EDTA	0.100
	glycerin	5.000
Phase B		
	cetyl hydroxy ethyl cellulose	0.150
Phase C		
	cetyl alcohol	0.500
	stearyl alcohol	0.500
	behenyl alcohol	0.500
	cetyl ricinoleate	3.000
	Steareth 2	1.000
	distearyl dimonium chloride	0.250
	myristyl myristate	1.500
	caprylic/capric/glyceride	1.500
	sucrose cottonate	1.000
Phase D		
	DI water	8.000
	niacinamide	2.000
Phase E		
	N,N"-methylene bis [N'- [(Hydroxymethyl)-2,5,-dioxy- 4-imidazolodiny] urea]	0.100
	DI water	0.300
	Isopropynyl butylcarbamide	0.005
	butylene glycol	0.300
Phase F		
	isopropyl palmitate	1.000
	retinol	0.050
	BHT	0.001
	Polysorbate 20	0.050

First, Phase A ingredients are combined and sparged using nitrogen for approximately 15 minutes. Phase B ingredients are then dispersed into Phase A until uniform using propeller type mixing and heated the mixture to about 75°C. In a separate vessel, Phase C ingredients are combined and heated to about 75°C. The mixture of phases A and B are then blanketed with a slow, steady stream of nitrogen.

10

- Next the Phase C ingredients are homogenized into the mixture of phases A and B using any rotor/stator type of homogenizer for approximately 15 minutes. After the 15 minutes, the mixing is switched to low rpm sweep mixing. Once the batch mixture is homogeneous, the entire batch mixture is cooled. When the batch is
- 5 cooled to about 50°C, phase D ingredients are added and homogenized. When the batch is cooled to about 40°C, phase E ingredients are added to the batch mixture. Lastly, when the batch mixture is cooled to about 30°C, the phase F ingredients are combined to the batch mixture. Mixing is continued until the batch mixture is uniform.
- 10 The resulting composition is useful for application to the skin for delivering the actives and to treat and improve the appearance of the skin.

## WHAT IS CLAIMED IS:

1. A skin care composition characterized in that it comprises:
  - a) from 0.01% to 50%, preferably from 0.1% to 10%, of a vitamin B<sub>3</sub> compound, and
  - b) from 0.001% to 5% of a preservative component, comprising:
    - i.) a formaldehyde donor; and
    - ii.) a halopropynyl compound selected from the group consisting of iodopropargyl esters, ethers, acetals, carbamates, carbonates and combinations thereof.
2. A composition according to Claim 1, wherein the weight ratio of the formaldehyde donor to halopropynyl compound is from 2000:1 to 1:1.
3. A composition according to Claim 1 or Claim 2, wherein the halopropynyl compound is 3-iodo-2-propynyl butyl carbamate.
4. A composition according to any of Claims 1 to 3, wherein the formaldehyde donor is selected from the group consisting of dimethyloldimethyldantoin, N,N"-methylene bis [N'-(hydroxymethyl)-2,5-dioxo-4-imidazolidinyl]urea; N-(hydroxymethyl)-N-(1,3-dihydroxymethyl-2,5-dioxo-4-imidazolidinyl)-N'-(hydroxymethyl) urea; the cis isomer of 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride sodium hydroxymethylglycinate, dimethyl oxazoline 7-ethylbicyclohexasolidine, 2-bromo-2-nitropropane-1,3-diol, 5-bromo-5-nitro-1,3-dioxane and mixtures thereof.
5. A composition according to any of Claims 1 to 4, wherein the vitamin B<sub>3</sub> compound is selected from the group consisting of niacinamide, derivatives of niacinamide, non-rubificent esters of nicotinic acid, and combinations thereof.
6. A composition according to any of Claims 1 to 5, wherein the vitamin B<sub>3</sub> compound is selected from the group consisting of niacinamide, tocopherol nicotinate, and combinations thereof.
7. A composition according to any of Claims 1 to 6, wherein said vitamin B<sub>3</sub> compound is substantially free of salts of the vitamin B<sub>3</sub> compound.

8. A composition according to any of Claims 1 to 7, wherein said vitamin B<sub>3</sub> compound is substantially uncomplexed.
9. A composition according to any of Claims 1 to 8, wherein said composition further comprises a compound selected from the group consisting of hydroxy acids, desquamatory agents, sunscreens, anti-oxidants, and combinations thereof.
10. A composition according to any of Claims 1 to 9 which further comprises from 1% to 99.5% of a dermatological carrier.
11. A composition according to any of Claims 1 to 10, wherein the carrier is of a form selected from the group consisting of solutions, aerosols, emulsions, lotions, creams, gels, sticks, sprays, ointments, pastes, mousses, makeup, and lipsticks.
12. A method of regulating skin conditioning, which is characterized in that it comprises applying to the skin of a mammal a safe and effective amount of a composition according to any of Claims 1 to 11.
13. A method of regulating visible and/or tactile discontinuities in the texture of mammalian skin, which is characterized in that it comprises applying to the skin of a mammal a safe and effective amount of a composition according to any of Claims 1 to 11.

# INTERNATIONAL SEARCH REPORT

In International Application No  
PCT/IB 98/00783

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A61K7/48 A61K7/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of the relevant passages	Relevant to claim No
A	US 5 254 331 A (MAUSNER JACK) 19 October 1993 see abstract see column 7, line 21 - line 41 see column 8, line 46 - line 59 see column 11, line 15 - line 33 see column 11, line 42 - line 49 ---	1, 5, 6, 9, 11-13
A	WO 97 17060 A (PROCTER & GAMBLE) 15 May 1997 see page 4, line 17 - page 5, line 26 see page 9, line 36 - page 10, line 7 see page 10, line 12 - line 15 see page 12, line 2 - line 9 --- -/--	1, 5-13

☒ Further documents are listed in the continuation of box C

☒ Patent family members are listed in annex

### Special categories of cited documents

- A\* document defining the general state of the art which is not considered to be of particular relevance
- E\* earlier document but published on or after the international filing date
- L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- O\* document referring to an oral disclosure, use, exhibition or other means
- P\* document published prior to the international filing date but later than the priority date claimed

- T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- &\* document member of the same patent family

Date of the actual completion of the international search

11 August 1998

Date of mailing of the international search report

19/08/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx 31 651 epo nl.  
Fax (+31-70) 340-3016

Authorized officer

McConnell, C

# INTERNATIONAL SEARCH REPORT

In International Application No  
PCT/IB 98/00783

C:(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	WO 81 00566 A (SUTTON LAB INC) 5 March 1981 see claim 1 see page 1, line 1 - page 4, line 24 ---	1,4
A	US 5 422 112 A (WILLIAMS LORRAINE) 6 June 1995 see column 5, line 35 - line 55 -----	1,3

# INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/IB 98/00783

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5254331	A	19-10-1993	NONE	
WO 9717060	A	15-05-1997	AU 7606196 A	29-05-1997
WO 8100566	A	05-03-1981	US 4271176 A	02-06-1981
			AT 11411 T	15-02-1985
			BE 906018 A	16-04-1987
			CA 1155129 A	11-10-1983
			CA 1236468 B	10-05-1988
			EP 0034182 A	26-08-1981
			JP 1054322 B	17-11-1989
			JP 56501047 T	30-07-1981
			US 4487939 A	11-12-1984
			US RE32848 E	31-01-1989
US 5422112	A	06-06-1995	CA 2150788 A	10-12-1995
			EP 0688560 A	27-12-1995
			JP 8053322 A	27-02-1996

